PMEL Tsunami Forecast Series: Vol. 11 A Tsunami Forecast Model for Point Reyes, California

Michael C. Spillane^{1,2}

¹Joint Institute for the Study of the Atmosphere and Ocean (JISAO), University of Washington, Seattle, WA

²NOAA/Pacific Marine Environmental Laboratory (PMEL), Seattle, WA

February 7, 2014

NOTICE from NOAA

Mention of a commercial company or product does not constitute an endorsement by NOAA/OAR. Use of information from this publication concerning proprietary products or the tests of such products for publicity or advertising purposes is not authorized. Any opinions, findings, and conclusions or recommendations expressed in this material are those of the author and do not necessarily reflect the views of the National Oceanic and Atmospheric Administration.



Contribution No. 3401 from NOAA/Pacific Marine Environmental Laboratory Contribution No. 2088 from Joint Institute for the Study of the Atmosphere and Ocean (JISAO)

Contents

Li	ist of Figures	
Li	ist of Tables	iii
1	Background and Objectives 1.1 The setting of Point Reyes, California	1 1 2 3
2	Forecast Methodology 2.1 The tsunami model	4 4 4
3	Model Development3.1 Digital elevation models	6 6 7 7 8 9
4	Results and Discussion 4.1 The "micro-tsunami" tests 4.2 The "mega-tsunami" tests 4.3 Model validation: the Honshu-2011 tsunami 4.4 Model validation with other preferred historical events 4.5 Other historical simulations of interest at Point Reyes, California 4.6 The Mendocino earthquake of April 25, 1992 4.7 Simulation of the remaining synthetic mega-events	11 11 12 14 16 18 19 20
5	Conclusions	22
6	Acknowledgments	23
7	References	24
FI	IGURES	54
T/	ABLES	98

Αŗ	Appendices	
A	Input files	107
	A.1 Reference model ★.in file for Point Reyes, California	108
	A.2 Forecast model ★.in file for Point Reyes, California	109
В	Propagation Database Pacific Ocean Unit Sources	110
C	SIFT Testing	150
	C.1 Purpose	150
	C.2 Testing procedure	150
	C.3 Results	151
Gl	Glossary	

List of Figures

1	The Point Reyes area of west and south Marin County, California	28
2	Extract from the oblique 3-D view of the San Francisco DEM provided by NGDC.	
	The focus is Point Reyes; areas of potential inundation identified by CalEMA are	
	highlighted in red.	29
3	View of the Point Reyes headland and Drake's Bay in its lee	30
4	Distribution of the historical tsunami sources employed for the development of	
	the Point Reyes forecast model. Those highlighted in red are more extensively	
	investigated using the reference model	31
5	A sample time interval from the Point Reyes tsunami-capable tide gauge, unre-	
	lated to tsunami activity. The evolving surface wave spectrum is shown in the	
	lower panel	32
6	The setting of Point Reyes and its nested forecast model grids. The C grids of other	
	west coast forecast models are marked, as are various sites with data available for	
	this study. The closest unit sources of the propagation database lie north of Cape	
	Mendocino, and the epicenter of the most recent Cascadia thrust event is marked.	33
7	Nested grid representation for the reference model (RM)	34
8	Nested grid representation for the forecast model (FM)	35
9	Comparison of the reference (RM) and forecast model (FM) time series at the	
	warning point for three "micro-tsunami" sources in the Western Pacific. The low-	
	est panel illustrates the appearance of model instability before the reference model	
	C grid bathymetry was finalized	36
10	Locations of synthetic tsunami scenarios employed in model development. Three	
	micro-tsunami scenarios, and the magnitude 7.5 case, employ a single unit source;	
	19 combine ten pairs of unit sources to model mega-tsunamis. Details are pro-	
	vided in Table ??. Cases highlighted in red have both forecast and reference model	
	solutions	37
11	Comparison of reference and forecast model results for the ACSZ 56-65 synthetic	
	mega-tsunami, representing the Cascadia Subduction Zone.	
	(a) distributions of maximum amplitude in the reference (upper panel) and fore-	
	cast model (lower panel) results with their time series (reference black, forecast	
	model red) and lagged correlation at the Point Reyes tide gauge as insets	38
11	(b) distributions of maximum speed in the reference (upper panel) and forecast	
	model (lower panel) results with the time series of the vector components at the	0.0
	Point Reyes tide gauge as insets	39

11	(c) comparison of the wave amplitude and currents in the reference (upper panel)	
	and forecast model (lower panel) results at the time indicated in the upper panel	40
	inset (the first wave peak)	40
11	(d) as in Figure ??c but at the later time when the reference and forecast model	41
10	solutions have diverged somewhat	41
12	As in Figure ?? but for the synthetic KISZ 01–10 event representing Kamchatka, (a) distributions of maximum amplitude in the reference (upper panel) and fore-	
	cast model (lower panel) results with their time series (reference black, forecast	
	model red) and lagged correlation at the Point Reyes tide gauge as insets	42
12	(b) distributions of maximum speed in the reference (upper panel) and forecast	42
12	model (lower panel) results with the time series of the vector components at the	
	Point Reyes tide gauge as insets	43
12	(c) comparison of the wave amplitude and currents in the reference (upper panel)	43
12	and forecast model (lower panel) results at the time indicated in the upper panel	
	inset (the first wave peak)	44
12	(d) as in Figure ?? c but for a time near the end of the simulation	45
13	As in Figure ?? but for the synthetic NTSZ 30–39 event representing Samoa,	
	(a) distributions of maximum amplitude in the reference (upper panel) and fore-	
	cast model (lower panel) results with their time series (reference black, forecast	
	model red) and lagged correlation at the Point Reyes tide gauge as insets	46
13	(b) distributions of maximum speed in the reference (upper panel) and forecast	
	model (lower panel) results with the time series of the vector components at the	
	Point Reyes tide gauge as insets	47
13	(c) comparison of the wave amplitude and currents in the reference (upper panel)	
	and forecast model (lower panel) results at the time indicated in the upper panel	
	inset (the first wave peak)	48
13	(d) as in Figure ??c but at the later time when the reference and forecast model	
	solutions have diverged somewhat	49
14	As in Figure ?? , but for a synthetic moderate event at NTSZ B36 near Samoa.	
	(a) distributions of maximum amplitude in the reference (upper panel) and fore-	
	cast model (lower panel) results with their time series (reference black, forecast	= 0
1.4	model red) and lagged correlation at the Point Reyes tide gauge as insets	50
14	(b) distributions of maximum speed in the reference (upper panel) and forecast	
	model (lower panel) results with the time series of the vector components at the	51
1.4	Point Reyes tide gauge as insets	51
14	and forecast model (lower panel) results at the time indicated in the upper panel	
	inset (the first major peak)	52
14	(d) as in Figure ??c but at the later time when the reference and forecast model	32
1-1	solutions have diverged somewhat.	53
15	Observed time series (red curves) from DART® and MARS bottom pressure sen-	00
	sors during the Honshu-2011 event, compared with the forecast model represen-	
	tation (black curves) based on the propagation database (see Table ?? ((a). Model	
	time series in the right hand panel have been lagged, and a common scale factor	
	of 1.2 applied	54

16	Comparison of observations with reference and forecast model-predicted time series at selected locations where tide gauge data are available: (a) Point Reyes, Arena Cove, and San Francisco. The time period highlighted in yellow marks an outage of 18 min or more that occured for the 1-min data streams along the entire west coast during the event. Runup values from the NGDC cat-	
	alog, when available, are indicated in the lower right of each panel for this and	
16	subsequent figures	55 56
17	Intercomparison of reference (upper panel) and forecast model (lower panel) predictions for the Honshu-2011 event.	30
	(a) maximum amplitude and reference point time series in the style of Figure ??a,	57
17	(b) maximum speed and reference point current time series in the style of Figure	
	?? b	58
17	(c) snapshot of amplitude and velocity at the leading wave peak in the style of	
	Figure ?? c,	59
17	(d) snapshot of amplitude and velocity at a later wave peak	60
17	(e) snapshot of amplitude and velocity at a later wave trough	61
18	Inundation forecast from the reference model (RM) C grid for the Honshu-2011	
	event, compared with the CalEMA inundation line. The inset in the upper right	
	shows tide gauge data from Point Reyes. Actual tides were well below MHW so	
	the inundation forecast was overly conservative	62
19	As in Figure 17 but for the Chile-2010 historical event,	
	(a) maximum amplitude and reference point time series in the style of Figure ??a,	63
19	(b) maximum speed and reference point current time series in the style of Figure	
• •	??b	64
19	(c) snapshot of amplitude and velocity at the leading wave peak in the style of	C.F.
20	Figure ??c	65 66
20 21	Model and observed time series comparison for the Chile-2010 event	66
21	(a) maximum amplitude and reference point time series,	67
21	(a) maximum amplitude and reference point time series,	68
21	(c) snapshot of amplitude and velocity at a time between waves at the reference	00
21	point	69
22	Model and observed time series comparison for the Samoa-2009 event	70
23	As in Figure 17 but for the Kuril-2006 historical event,	10
_0	(a) maximum amplitude and reference point time series,	71
23	(b) maximum speed and reference point current time series,	72
23	(c) snapshot of amplitude and velocity at the leading wave peak	73
24	Model and observed time series comparison for the Kuril-2006 event	74
25	As in Figure 17 but for the Alaska-1964 historical event,	
	(a) maximum amplitude and reference point time series,	75
25	(b) maximum speed and reference point current time series,	76
25	(c) snapshot of amplitude and velocity at the leading wave peak	77
26	Model and observed time series comparison for the Alaska-1964 event	78
27	As in Figure 17 but for the Unimak-1946 historical event,	
	(a) maximum amplitude and reference point time series,	79

27	(b) maximum speed and reference point current time series,	80
27	(c) snapshot of amplitude and velocity at the later wave peak	81
28	Model and observed time series comparison for the Unimak-1946 event	82
29	The Sanriku event of June 15, 1896	83
30	The Kamchatka event of November 4, 1952	84
31	The Chile event of May 22, 1960	85
32	The Andreanof event of June 10, 1996	86
33	The Peru event of June 23, 2001	87
34	The Hokkaido event of September 25, 2003	88
35	The Rat Island event of November 17, 2003	89
36	The Tonga event of May 3, 2006	90
37	The normal thrust event off the Kuril Islands on January 13, 2007	91
38	The Solomon event of April 1, 2007	92
39	The Peru event of August 15, 2007	93
40	The Chile event of November 14, 2007.	94
41	The Cape Mendocino event of April 25, 1992. The upper panels show the fre-	
	quency of non-thrust events in the vicinity, with only two having a focal mecha-	
	nism characteristic of subduction. Lower panel: comparison of model with ob-	
	servation at Arena Cove and Point Reyes	95
42	Predicted maximum sea level (from the forecast model) at the Point Reyes tide	
	gauge that might result were "mega-tsunamis" to occur at various locations around	
	the Pacific basin.	96
43	Chart of the area inundated by one or more of the mega-tsunami scenarios based	
	on the forecast model. Shown in blue is the CalEMA inundation line, which is	
	based on a similar ensemble of scenarios	97
		111
B.2		117
B.3		125
B.4	Kamchatka-Bering Subduction Zone unit sources	
B.5		129
	J J	134
		136
		138
		142
	New Zealand–Puysegur Zone unit sources	146
B.11	Ryukyu–Kyushu–Nankai Zone unit sources	148
C.1	Response of the Point Reyes forecast model to synthetic scenario KISZ 1-10 (α =	
	25). Maximum sea surface elevation for (a) A grid, (b) B grid, (c) C grid. Sea sur-	
	face elevation time series at the C grid warning point (d). The extrema at the	
	reference point are compared with the equivalent values obtained during model	
		153

C.2	Response of the Point Reyes forecast model to synthetic scenario KISZ 22-31 (α =	
	25). Maximum sea surface elevation for (a) A grid, (b) B grid, (c) C grid. Sea sur-	
	face elevation time series at the C grid warning point (d). The extrema at the	
	reference point are compared with the equivalent values obtained during model	
	development listed in Table C.1	154
C.3	Response of the Point Reyes forecast model to synthetic scenario ACSZ 56-65 (α =25)	
	Maximum sea surface elevation for (a) A grid, (b) B grid, (c) C grid. Sea surface	
	elevation time series at the C grid warning point (d). Panels (c) and (d) can be	
	compared with the equivalent results, obtained during model development, dis-	
	played in Figure ??	155
C.4	Response of the Point Reyes forecast model to synthetic scenario CSSZ 89-98 (α =	
	25). Maximum sea surface elevation for (a) A grid, (b) B grid, (c) C grid. Sea sur-	
	face elevation time series at the C grid warning point (d). Panels (c) and (d) can	
	be compared with the equivalent results, obtained during model development,	
	listed in Table C.1	156
C.5	Response of the Point Reyes forecast model to synthetic scenario NTSZ 30-39	
	($\alpha = 25$). Maximum sea surface elevation for (a) A grid, (b) B grid, (c) C grid.	
	Sea surface elevation time series at the C grid warning point (d). Panels (c) and	
	(d) can be compared with the equivalent results, obtained during model develop-	
	ment, displayed in Figure ??	157
C.6	Response of the Point Reyes forecast model to the March 11 Tohoku tsunami. (a,	
	b, and c) Maximum sea surface elevation for A, B and C grids. (d) Sea surface	
	elevation time series at the C grid warning point. Panels (c) and (d) can be com-	
	pared with the equivalent results, obtained during model development, displayed	
	in Figure ?? and Figure ?? a	158

List of Tables

1	Source characterization for historical tsunami events employed in Point Reyes model testing. Those in bold text were used to compare the reference and fore-	
	cast model versions. Sources identified as "preliminary" or "ad hoc" may not be	
	identically defined in other forecast model reports.	
1	a) The standard set for Pacific Ocean models	99
	forecast model testing.	100
2	The main features of the San Francisco digital elevation model (DEM), which in-	
	cludes Point Reyes, California	101
3	Tidal characteristics of the Point Reyes, California tide gauge (9415020)	102
4	Specifics of the reference (RM) and forecast model (FM) grids employed for Point	
	Reyes, California. For the paired values in the resolution and grid points columns,	
	the zonal (East to West) value is listed first, followed by meridional (North to	
	South). CPU times for a 4-hr simulation are based on use of a single Intel [®] Xeon [®]	
	E5670 2.93GHz processor	103
5	Grid file names and grid-related parameters. The time steps for the A and B grids	
	must be integer multiples of the basic time step chosen for the C grid	104
6	Synthetic tsunami events employed in Point Reyes, California model testing. The	
	reference and forecast model solutions of those shown in bold text were inter-	
	compared extensively	105
7	Mega-tsunami scenario impacts, represented by flooding (km² in the sub-region	
	123.1°-122.5°W, 37.8°-38.1°N) and maximum amplitude (cm) at several sites within	
	the model domain (identified in the footnote). The maxima are highlighted and	
	ranked	106
B.1	Earthquake parameters for Aleutian-Alaska-Cascadia Subduction Zone unit sources	s.112
B.2	Earthquake parameters for Central and South America Subduction Zone unit source	es.118
B.3	Earthquake parameters for Eastern Philippines Subduction Zone unit sources	126
B.4	Earthquake parameters for Kamchatka-Bering Subduction Zone unit sources	128
B.5	Earthquake parameters for Kamchatka-Kuril-Japan-Izu-Mariana-Yap Subduction	
	Zone unit sources	130
B.6	Earthquake parameters for Manus–Oceanic Convergent Boundary unit sources	135
B.7	Earthquake parameters for New Guinea Subduction Zone unit sources	137
B.8	Earthquake parameters for New Zealand–Kermadec–Tonga Subduction Zone unit	
	sources	139

B.9	Earthquake parameters for New Britain–Solomons–Vanuatu Subduction Zone unit	
	sources	143
B.10	Earthquake parameters for New Zealand–Puysegur Subduction Zone unit sources.	147
B.11	Earthquake parameters for Ryukyu–Kyushu–Nankai Subduction Zone unit sources.	149
C.1	Table of maximum and minimum amplitudes (cm) at the Point Reyes, California warning point for synthetic and historical events tested using SIFT 3.2 and those	
	obtained during development	159



Foreword

Tsunamis have been recognized as a potential hazard to United States coastal communities since the mid-twentieth century, when multiple destructive tsunamis caused damage to the states of Hawaii, Alaska, California, Oregon, and Washington. In response to these events, the United States, under the auspices of the National Oceanic and Atmospheric Administration (NOAA), established the Pacific and Alaska Tsunami Warning Centers, dedicated to protecting United States interests from the threat posed by tsunamis. NOAA also created a tsunami research program at the Pacific Marine Environmental Laboratory (PMEL) to develop improved warning products.

The scale of destruction and unprecedented loss of life following the December 2004 Sumatra tsunami served as the catalyst to refocus efforts in the United States on reducing tsunami vulnerability of coastal communities, and on 20 December 2006, the United States Congress passed the "Tsunami Warning and Education Act" under which education and warning activities were thereafter specified and mandated. A "tsunami forecasting capability based on models and measurements, including tsunami inundation models and maps..." is a central component for the protection of United States coastlines from the threat posed by tsunamis. The forecasting capability for each community described in the PMEL Tsunami Forecast Series is the result of collaboration between the National Oceanic and Atmospheric Administration office of Oceanic and Atmospheric Research, National Weather Service, National Ocean Service, National Environmental Satellite, Data, and Information Service, the University of Washington's Joint Institute for the Study of the Atmosphere and Ocean, National Science Foundation, and United States Geological Survey.

PMEL Tsunami Forecast Series: Vol. 11

A Tsunami Forecast Model for Point Reyes, California

M. C. Spillane ¹, ²

Abstract. Operational tsunami forecasting by NOAA's Tsunami Warning Centers relies on the detection of tsunami wave trains in the open ocean, inversion of these data (telemetered via satellite) to quantify their source characteristics, and real-time modeling of the impact on threatened coastal communities. The latter phase of the process involves, for each such community, a pre-tested forecast model capable of predicting the impact, in terms of inundation and dangerous inshore currents, with sufficient resolution and within the time constraints appropriate to an emergency response.

In order to achieve this goal, considerable advance effort is required to tune each forecast model to the specific bathymetry and topography, both natural and manmade, of the impact area, and to validate its performance with a broad set of tsunami sources. Where possible the validation runs should replicate observed responses to historical events, but the sparse instrumental record of these rare but occasionally devastating occurrences dictates that comprehensive testing should also include a suite of scenarios that represent potential future events.

During the forecast model design phase, and in research mode outside the pressures of an emergency situation, more detailed and slower-running models can be investigated. Such a model, referred to as a reference model, represents the most credible numerical representation of tsunami response for the study region, using the most detailed bathymetry available and without the run-time constraint of operational use. Once a reference model has been developed, the process of forecast model design is to determine where efficiencies can be gained, through reducing the grid resolution and increasing the model time step, while still adequately representing the salient features of the full solution.

This report documents the reference and forecast model development for Point Reyes and vicinity, comprising much of western Marin County, California. The Point Reyes headland juts out into the Pacific and its lighthouse is a prominent navigation landmark northwest of the entrance to San Francisco Bay. A tide gauge within Drake's Bay, in the lee of the headland, provides observations for model validation from numerous historical tsunamis. While much of the study region lies within a National Seashore area, limiting the population and waterfront infrastructure, there are a number of nearby communities exposed to tsunami impact. Beaches and other natural amenities and the mild climate foster extensive recreational use, and there is a clear need for emergency preparedness. This report addresses the tsunami aspects of the natural hazard spectrum.

 $^{^{1}}$ Joint Institute for the Study of the Atmosphere and Ocean (JISAO), University of Washington, Seattle, WA

²NOAA/Pacific Marine Environmental Laboratory (PMEL), Seattle, WA

Background and Objectives

1.1 The setting of Point Reyes, California

Point Reyes, lying to the northwest of the entrance to San Francisco Bay, is a prominent navigational landmark. As illustrated in Figure 1, composed of orthographic images from "Marin-Maps" (mmgis.marinmap.org/OrthoGrid/viewer.htm), the headland is the site of a lighthouse and, in Drake's Bay in its lee adjacent to the historic Point Reyes Lifeboat station, is the tide gauge bearing the same name. All lie within the Point Reyes National Seashore (PRNS), comprising most of west Marin County, which is essentially unpopulated and in a natural state, with the exception of some agricultural activity that was allowed to continue when the PRNS was established in 1962. As seen in the inset to Figure 1, the San Andreas Fault strongly delineates the eastern boundary of the region though it is submerged in Tomales Bay, in the north, and Bolinas Lagoon in the south. In the neck of land between them are the communities of Olema and Point Reyes Station, which are close to the epicenter of the 1906 San Francisco Earthquake. Several small communities lie on the shores of Tomales Bay which is 20.4 km in length but with a mean depth of only 3.1 m, (Niemi and Hall, 1996). The entrance is shallow and constricted. At the southern end, Bolinas (2010 Pop. 1620) and Stinson Beach (2010 Pop. 632) have greater exposure to damage from tsunami or winter storm waves. Between Stinson Beach and Point Bonita, the southernmost point of Marin County, lies Muir Beach, a community of about 310. It is notable, from the tsunami perspective, in that it reported major run-ups during the Unimak-1946 and Alaska-1964 events.

North of Tomales Bay is Bodega Bay whose shores lie both in Marin and Sonoma counties. Apart from the shallow Bodega Harbor, and the communities of Bodega Bay (2010 Pop. 1077) and Doran Beach extending onto the spit at its mouth, this area too is sparsely populated. The natural beauty of the region, with its mild climate and proximity to the San Francisco area and other urban centers, provides outstanding recreational opportunities, resulting in large numbers of visitors throughout the year. Normally, in selecting the domain of a tsunami forecast model, the location of a tide gauge provides the focus but, in this case, a somewhat larger region is included to provide forecast capability to population centers and primary recreational assets. Initially it was hoped that a forecast model could cover the entire region from Bodega Bay to Muir Beach. This proved to be impossible, given the time constraints on model runtime imposed by emergency usage, without an unacceptable reduction in spatial resolution. While the innermost of the nested grids in the reference model does include Tomales Bay and a portion of Bodega Bay, these are excluded from the forecast model, which focuses on the

south and southwest of Marin County. http://xenon.colorado.edu/spotlight/ The University of Southern California Tsunami Research Center conducted a comprehensive study of potential tsunami inundation, for the entire California coastline. Funded through the California Emergency Management Agency (CalEMA), by the National Tsunami Hazard Program, the study (Barberopoulou et al., 2011) has produced a set of inundation maps for emergency planning purposes accessible online in various forms, including MyHazard (myhazards.calema.ca.gov) which enables users to acquire information specific to their site of interest. The CalEMA inundation results are available in GIS form and those specific to the Point Reyes area are used throughout this report. In addition to underpinning the modeling effort, the digital elevation model (DEM) for the San Francisco region, provided by the National Geophysical Data Center (NGDC), includes a 3-D oblique view that assists greatly in visualizing the study area. In Figure 2, the CalEMA inundation information is overlaid, together with descriptive labels on an extract from the NGDC image. The full 3-D image is available in the San Francisco DEM Report (Carignan et al., 2010).

A striking series of aerial photographs (www.californiacoastline.org) show that the study region contains both high cliffs (also seen in Figure 3) that limit potential impact by tsunamis and broad beaches and shallow coastal inlets that are more exposed. Queries to the CalEMA "My-Hazards" site for Bolinas and Stinson Beach show flooding and earthquake as other hazards to which they are prone, in addition to tsunami. Available online is a video "Marin Tsunami" (Loeffler and Gesell, 2010), produced for the United States Geological Survey (USGS) in cooperation with the Marin County Sheriff's Office of Emergency Services. In addition to providing an overview of the comprehensive level of preparedness for tsunami impact on the communities of Bolinas, Stinson Beach, Dillon Beach / Lawson's Landing, and the popular Limantour Beach within the National Parks area of responsibility, this excellent resource for residents and visitors alike gives insight into the character of the area.

1.2 Natural hazards

Instances of mild tsunami signals are evident in the tide gauge records for Point Reyes (established in 1975), and Marin County sites appear several times in the records compiled by Lander and Lockridge (1989) and its regularly updated online equivalent: NGDC Tsunami Hazard Database (Dunbar (2007); see ngdc.noaa.gov/hazard/). The historical record first mentions Marin County with a wave observed at Sausalito, on the north shore of the Golden Gate, from a Chilean event in 1877. The earliest time series currently available for analysis is a digitized marigram from Sausalito, recorded during the Sanriku event of 1896 and available in the Alaska / West Coast Tsunami Warning Center (WCATWC) archives. O'Brien (1946) described a 2.6 m wave (above MLLW) in Drake's Bay during the 1946 Unimak tsunami, with a boat washed onto the highway. While Marin County sites are not explicitly mentioned in connection with the Kamchatka-1952 or Andreanof-1957 events, waves were observed at Bodega and within San Francisco Bay. During the 1960 Chile event, a 1.5 m run-up was reported at Stinson Beach, and during the 1964 Alaska tsunami waves were observed at several sites within Marin County including Drake's Beach. Time series from several tsunamis are available from the Point Reyes tide gauge in recent years, culminating in the major event east of Honshu on March 11, 2011 (also referred to as the Tohoku earthquake.) The latter will be discussed extensively in this report.

Combining events impacting northern California with those that have occurred since the

Point Reyes tide gauge was upgraded to 1-min sampling, a total of 27 historical events are available for study. Nineteen of these, listed in Table 1a, are among the preferred cases for forecast model testing in the Pacific because their sea floor deformation is reasonably well known, either from the literature or, more recently, derived from direct observation of the wave trains they generated. The remaining eight, listed in Table 1b, have source characteristics that are less well known; they are included to expand the geographical coverage or because of their special relevance to the west coast. The Mw 7.2 earthquake north of Cape Mendocino on April 25, 1992 was a very mild foretaste of a Cascadia Subduction Zone (CSZ) event but was registered in marigrams at Arena Cove and Point Reyes. Others, due to significant noise in the tide gauge, do not produce a clear signal but shed light on Point Reyes as a reference point for coastal impacts. Figure 4 illustrates the distribution of the 27 historical sources. Those highlighted in red were employed for intercomparison of the reference and forecast versions of the model.

Direct seismic impact is another natural hazard to which Marin County is exposed. Its proximity to the rupture zone of the SAF in the San Francisco earthquake of 1906 resulted in significant lateral displacements and some damage in the inland towns. The lighthouse on the Point Reyes headland suffered only mild damage. While the SAF enters the ocean at Bolinas, its strike-slip nature reduces the likelihood of severe tsunami wave generation should ruptures occur in the immediate vicinity. Submarine landslides or collapse of sections of sea cliff are, however, a potential local source for tsunami damage. Landslides triggered by seismic events caused significant loss of life during the 1929 Newfoundland event and accentuated the 1996 New Guinea tsunami. Landslide-generated tsunami waves are not currently included in the forecasting system SIFT (Short-term Inundation Forecasting for Tsunamis), developed at NCTR (NOAA Center for Tsunami Research) and now in operational use at the U.S. Tsunami Warning Centers (TWC), nor are those generated meteorologically. However, to the extent that the waves they produce are detected by the DART® array, some warning of their presence may be available.

Another local hazard that has been a frequent cause of damage in the Bolinas-Stinson Beach area is ocean wave action. Originating locally, or as swell from distant storms, such waves in the winters of 1977-78 and 1982-83 caused the loss of several beachfront homes. Another impact of ocean waves, of relevance to tsunami detection and modeling, is in the noise they produce in the tide gauge records. Although the Point Reyes tide gauge is in the lee of the headland, excessive wave action and resonance can mask weak tsunami signals.

1.3 Tsunami warning and risk assessment

The forecast model development, described here, will permit Point Reyes, California, to be incorporated into the tsunami forecasting system SIFT. The system has had considerable success in accurately forecasting the impact of both moderate and severe tsunami events in recent years and in the following section the methodology that permits such forecasts is discussed as prelude to a description of forecast model development for Point Reyes. With the model in hand, validated with historical events and with its stability verified by extensive testing against extreme scenarios, real-time forecasts will be available to inform local emergency response. Additionally, the synthetic scenarios investigated during model development, and reported here, provide an initial tsunami risk assessment as described in the Results and Discussion section.

Forecast Methodology

2.1 The tsunami model

In operational use, NOAA's tsunami forecast model is used to extend a pre-computed deepwater solution into the shallows, and onshore as inundation if appropriate. The model consists of a set of three nested grids named A (outermost with coarse resolution), B (intermediate), and C (innermost). The latter provides fine resolution that, in a real-time application of the MOST model (Method of Splitting Tsunami: Titov and Synolakis (1998); Titov and González (1997)), permits forecasts at spatial scales (as little as few tens of meters) relevant to local emergency management. The validity of the MOST model applied in this manner, and the operational effectiveness of the forecast system built around it, has been demonstrated during unplanned tests in the Pacific basin triggered by several mild to moderate tsunami events in the years since the 2004 Indian Ocean disaster (Wei et al., 2008), and during the severe Honshu-2011 tsunami. Successful hindcasting of observed historic events, even mild ones, during forecast model development lends credence to an ability to accurately forecasting the impact of future events. Such validation of tsunami modeling procedures is documented in other volumes of the series of which this report is but one. Before proceeding to a description of the forecast model development for Point Reyes, California, it is useful to describe the steps in the overall forecast process.

2.2 NOAA's tsunami forecast system

Operational tsunami forecasts are generated at Tsunami Warning Centers, staffed 24/7 in Alaska and Hawaii, using the SIFT tool, developed at NCTR. The semi-automated process facilitates the steps by which TWC operators assimilate data from an appropriate subset of DART® tsunami sensors, "invert" the data to determine the linear combination of pre-computed propagation solutions that best match the observations, then initiate a set of forecast model runs if coastal communities are threatened or, if warranted, cancel the warning. Steps in the process are as follows:

• When a submarine earthquake occurs the global network of seismometers registers it. Based on the epicenter, the unit sources in the Propagation Database (Gica et al., 2008) that are most likely to be involved in the event, and the DART® array elements (Spillane et al., 2008) best placed to detect the waves passage are identified. TWC operators can

trigger DART®s into rapid sampling mode in the event that this did not occur automatically in response to the seismic signal.

- There is now an unavoidable delay while the tsunami waves are in transit to the DART[®]s; at least a quarter of a cycle of the first wave in the train must be sampled before moving to the "inversion" step.
- When sufficient data have accumulated, at one or more DART[®]s, the observed time series are compared with the model series from the candidate unit sources. Since the latter are pre-computed (using the MOST code), and the dynamics of tsunami waves in deep water is linear, a least squares approach taking very little time can identify the unit sources (and the appropriate scale factors for each), that best fit the observations. The "inversion" methodology is described by Percival et al. (2011).
- Drawing again on the Propagation Database, the scale factors are applied to produce a composite basin-wide solution with which to identify the coastal regions most threatened by the radiating waves.
- It is at this point that one or more forecast models are run. The composite propagation solution is employed as the boundary condition to the outermost (A grid) domain of a nested set of three real-time MOST model grids that telescope with increasingly fine scale to the community of concern. A grid results provide boundary conditions to the B grid, which in turn forces the innermost C grid. Non-linear processes including inundation are modeled so that, relying on the validation procedures during model development, credible forecasts of the current event are available.
- Each forecast model provides quantitative and graphic forecast products with which to inform the emergency response, or to serve as the basis for canceling or reducing the warnings. Unless the tsunami source is local, the forecast is generally available before the waves arrive but, even when lead-time cannot be provided, the several hour duration of a significant event (in which the first wave may not be the most damaging) gives added value to the multi-hour forecasts provided.

Because multiple communities may be potentially at risk, it may be necessary to run simultaneously, or in a prioritized manner, multiple forecast models. Each must be optimized to run efficiently in as little time as possible; the current target is that an operational forecast model should be capable of simulating 4 hr of real time within about 10 min of CPU time on a fast workstation computer.

Model Development

3.1 Digital elevation models

Water depth determines local tsunami wave speed and sub-aerial topography determines the extent to which tsunami waves inundate the land. Thus a prerequisite for credible tsunami modeling is the availability of accurate gridded bathymetric and topographic datasets, termed digital elevation models (DEM). Given their expertise in this area, and the number of coastal communities needing tsunami forecast capability, NCTR relies heavily on the National Geophysical Data Center (NGDC) to provide the DEMs needed. In the case of Point Reyes, California, a sub-region of the San Francisco DEM is employed. This DEM, a composite of multiple data sources merged and converted to a common datum of Mean High Water (MHW), was produced and documented by Carignan et al. (2010). MHW as the "zero level" is employed in all forecast models. The MOST model does not include tidal fluctuations and, since a tsunami may arrive at any stage of the tide, it is best to employ a "worst-case" approach by assuming high tide when forecasting inundation. For some forecast models grounding of vessels and the strong and rapidly varying currents often associated with even mild tsunamis are of concern. For Point Reyes, lacking a marina and shoreline infrastructure, low water impacts are less important.

The Point Reyes sub-region of the San Francisco DEM was illustrated in Figure 2; its salient features listed in Table 2 are reproduced from DEM documentation (Carignan et al., 2010). The NGDC report thoroughly describes the data sources and methods employed in constructing the DEM. With one-third arcsec (10 m) resolution, the DEM provides the basis for the B and C grids for both reference and forecast model usage. NCTR maintains an atlas of lower resolution gridded bathymetries, which can be used for the A grids, as described later. All of the DEMs employed were verified for consistency with charts, satellite imagery, and other datasets during the course of MOST grid development.

The elevations and depths used in the development of this forecast model were based on the digital elevation model provided by the National Geophysical Data Center and the author considers it to be a good representation of the local topography and bathymetry. As new digital elevation models become available, forecast models will be updated and report updates will be posted at nctr.pmel.noaa.gov/forecast_reports/.

3.2 Tides and sea level variation

The history of tidal observation at Point Reyes, California dates back only to 1975. The tide station (9415020) is located near the end of a pier projecting into Drake's Bay just west of the historic Lifeboat station. The pilings raise the deck well above sea level and do not impede water movement. The instrumentation was upgraded in 2006 to include a tsunami-capable gauge sampling at 1-min intervals (and, on demand, at 15-sec intervals); some earlier data were sampled at 6-min intervals and several historical events are only available as marigrams on microfiche. An ongoing project at NGDC will digitize the more critical images in this archive; a few are available in digitized form in the WCATWC archives.

Station characteristics for 9415020 are provided in Table 3, based on the wealth of online tidal information available at NOAA's Center for Operational Oceanographic Products and Services (CO-OPS) website (tidesandcurrents.noaa.gov). Note the sizeable diurnal range of about 1.7 m and that, while the long-term rate of change in sea level is low (compared to more tectonically active areas), there is substantial seasonal, interannual and short-term variability.

A sample section of the tide gauge record, extracted from the CO-OPS website for the period following the Honshu-2011 is included in Figure 18 from Section 4.3. In a several hour section of 1-min data, the signature of an arriving tsunami is generally a burst of higher frequency energy with a sudden onset. However, during the winter months in particular, similar bursts unrelated to tsunami activity are quite common. In January 2011 for example several occurred, one of which is illustrated in Figure 5. The tidal signal has been removed with a Butterworth band-pass filter with cutoff periods at 5 and 120 min. (This filter is used throughout the report to preprocess tide gauge records for comparison with model prediction.) The lower panel of Figure 5 is the spectral wave energy at hourly intervals from NDBC buoy 46026, 18 nm west of San Francisco. There is a clear correlation between enhanced swell at this site and the detided residuals in Drake's Bay suggesting that surface waves can excite a coastal response. The amplitude of this noise for the example shown is perhaps 10-20 cm and would likely obscure a mild tsunami's signature were one to arrive during such an episode. Deviations (or residuals) from the astronomically predicted tide can be several centimeters and the variability strong. In particular the highest water level reported for the Point Reyes gauge is 1.05 m above MHW (Feb 6, 1998) so the use of MHW as the zero level of modeled sea level may underestimate the truly worse case. While the simultaneous arrival of the crest of a large tsunami at high tide during a storm surge has low probability, a feature of the simulated events reported below is that sustained oscillations at a resonant period may extend the duration of the threat. This effect is notorious at Crescent City, California which is frequently the most heavily impacted U.S. west coast location for remote events.

3.3 The CFL condition and other considerations for grid design

Water depth dependent wave speed, in conjunction with the spacing of the spatial grid representation, places an upper limit on the time step permissible for stable numerical solutions employing an explicit scheme. This is the CFL (Courant-Friedrichs-Levy) limit, which requires careful consideration when the grids employed for a reference or forecast model are being designed. Finer-scale spatial grids, or greater water depths, require shorter time steps thereby increasing the amount of computation required to simulate a specific real time interval.

Another feature of the application of gridded numerical solutions to the tsunami wave

problem is the shortening that the wave train encounters in moving from deep water onto the shelf. In deep water a grid spacing of 4 arcsec (of latitude and longitude, corresponding to \sim 7 km) is normally used to represent propagating wave trains whose wavelength is typically of the order of a few hundred kilometers. The stored results of such propagation model runs are typically decimated by a factor of 4, resulting in a database of \sim 30 km spacing (and 1-min temporal sampling) with which to generate the boundary conditions for the outermost of the nested grids in a model solution. The extraction of the boundary conditions (of wave height and the two horizontal velocity components) is achieved by linear interpolation in space and time. To provide realistic interpolated values the stored fields for these variables must be smoothly varying, and have adequate sampling in space and time to resolve their structure. This necessitates the placement of the offshore boundary of the forecast model domain well offshore. The presence of the Mendocino Escarpment is another incentive to do so, in order that its role in topographic steering of trans-Pacific wave trains is adequately represented.

3.4 Specifics of the model grids

After several rounds of experimentation, the extents and resolutions of the nested grids were chosen, and are illustrated in Figures 6, 7 and 8; details are provided in Tables 4 and 5. The reference model grid extents were set early in the process when the hope was to provide forecast results from Bodega Bay to Muir Beach, but have further value in ensuring adequate representation of waves entering the domain from remote sources. The reference model grids are displayed in Figure 7; in the A and B grid panels, rectangles show the nested grid domain within. In the case of the reference model C grid panel, the reduced extent of the equivalent forecast model grid is indicated. Figure 8 depicts the nested grids of the forecast model itself. The main focus of the forecast model, and this report, is on the southwestern and southern portion of Marin County. Some mention of the northern portion will be made as appropriate but, with the exception of some results that can be derived from the A grid, comprehensive forecasts for Bodega Bay will require a dedicated model.

Both C grids lie entirely within the NGDC-provided DEM; A and B grids include bathymetry and topography from other DEM datasets available at NCTR. Some smoothing and editing were necessary to eliminate erroneous points or grid features that tend to cause model instability. For example, "point" islands where an isolated grid cell stands above water are eliminated, as are narrow channels or inlets one grid unit wide; these tend to resonate in the numerical solution. Large depth changes between adjacent grid cells can also cause numerical problems; customized tools (such as "bathcorr") are available to correct many of these grid defects. An additional constraint on the bathymetry (Elena Tolkova, personal communication), which identifies excessive depth changes in the discrete representation, was applied.

Details of the model grids are provided in Tables 4 and 5. The latter lists the maximum depth, the CFL time step requirement that must not be exceeded, and the actual time steps chosen for the reference and forecast model runs. Since, in the current version of MOST employed by SIFT, the numerical solutions in the three grids proceed simultaneously, there is a requirement that the A and B grid time steps be integer multiples of the (innermost) C grid time step in addition to satisfying the appropriate CFL requirement. For both reference and forecast models the CFL requirement of the C grid was the most stringent. The values chosen are shown in the final column of Table 4 and are such that an integer multiple of each time step $(15 \times \text{ for the forecast model}; 50 \times \text{ for the reference})$ is identically 30 sec, the chosen output time

interval for both models. When run on an Intel[®] Xeon[®] E5670 2.93GHz processor the forecast model produces 4 hr of simulation in 7.46 min, within the desired 10-min value for this metric.

3.5 Model run input and output files

In addition to providing the model grid file names and the appropriate time step and A, B grid multiples as provided in the tables above, it is necessary to provide a number of additional parameters in an input file. These include the Manning friction coefficient, a depth threshold to determine when a grid point becomes inundated, and the threshold amplitude at the A grid boundary that will start the model. An upper limit on wave amplitude is specified in order to terminate the run if the waves grow beyond reasonable expectation. Usual MOST values are used: 0.0009 for the friction coefficient and 0.1 m for the inundation threshold. The latter causes the inundation calculation to be avoided for insignificant water encroachments that are probably below the level of uncertainty in the topographic data. Inundation can, optionally, be ignored in the A and B grids, as is the norm in the (non-nested) MOST model runs that generate the propagation database. When A and B grid inundation is excluded, water depths less than a specified "minimum offshore depth" are treated as land; in effect a "wall" is placed at the corresponding isobath. When invoked, a typical value of 1-5 m is applied as the threshold, though A and B inundation is normally permitted as a way to gain some knowledge of tsunami impact beyond the scope of the C grid domain. Other parameter settings allow decimation of the output in space and/or time. As noted earlier, 30-sec output has been the target and output at every spatial node is preferred. These choices avoid aliasing in the output fields that may be suggestive of instability (particularly in graphical output), when none in fact exists.

Finally the input file (supplied in Appendix A) provides options that control the output produced. Output of the three variables: wave amplitude, and the zonal (positive to the east) and meridional (positive to the north) velocity components can be written (in netCDF format) for any combination of the A, B, and C grids. These files can be very large! A separate file, referred to as a "SIFT" file, contains the time series of wave amplitude at each time step at discrete cells of a selected grid. Normally the time series at a "reference" or "warning point", typically the location of a tide gauge, is selected to permit validation in the case of future or historical events. As noted earlier, several additional sites within the model domain were specified during development and are included in subsequent discussion in this report. Also output in the SIFT file is the distribution of the overall minimum and maximum wave amplitude and speed in each grid. By contrast with the complete space-time results of a run, the SIFT file (also netCDF) is very compact.

By default two additional output files are generated. A listing file summarizes run specifications, progress, and performance in terms of run-time. Also included in this file is information to determine the reason, should a run not start or terminate early. Finally a "restart" file is produced so that a run can be resumed, beginning at the time it ended, either normally or by operator intervention.

The input files described above are specific to the model itself. For an actual run, the program must be pointed toward the files that contain the boundary conditions of wave amplitude (HA), and velocity components (UA, VA), to be imposed at the A grid boundary. Time varying conditions are generally extracted as a subset of a basin-wide propagation solution (either a single unit source or several, individually scaled and linearly combined) that mimic a particular event. These boundary-forcing files typically consist of 24 hr of values (beginning at the

time of the earthquake), sampled at 1-min intervals and available on a 16 arcmin grid. Occasionally, for more remote seismic sources (or when delayed arrival of secondary waves due to reflections are a concern, as has been seen at Hawaii, the time span of the propagation run available for forcing is extended beyond one day.



Results and Discussion

Before proceeding to an extensive suite of model runs, that explore the threat to Marin County from various source regions, the stability of the model is tested in both low and extreme amplitude situations. The former we refer to as "micro-tsunami" testing: where the boundary forcing is at such a low level (but not precisely zero of course) that the response is expected to be negligible. These tests can be highly valuable in revealing localized instabilities that may result from undesirable features in the discretized bathymetric representation. Inlets or channels that are only one grid cell wide may "ring" or resonate in a non-physical way in the numerical solution. An instability may not grow large enough to cause the model to fail but, in a run with typical tsunami amplitudes, may be masked by actual wave variability.

Forcing by extreme events should also be tested. In addition to the need to test model stability under such circumstances, there is a parameter in the input file that truncates the run if a prescribed threshold is exceeded. For operational use, the threshold must be set high enough so that an extreme event run is not unnecessarily terminated. Both tests should be performed for test sources whose waves enter the model domain from different directions since, although stable for one set of incoming waves, an instability may be encountered for another. The "micro-" and "mega-tsunami" testing of the forecast and reference models is reported in the following subsections. Further evidence of stability is provided by the extensive set of scenarios, aimed at exploring the dependence of impact to source location, described later in the report, and in independent testing by other members of the NCTR team before the model was released for operational use.

4.1 The "micro-tsunami" tests

Three cases (see Table 6) were run representing micro-tsunami sources in the western Aleutians, the Philippines, and south of Japan (see Figure 10). Based on sources from the propagation database (Gica et al., 2008), their amplitudes were scaled down by a factor of 100 so as to mimic an Mw 6.167 / Slip 0.01 m source rather than the Mw 7.5 / Slip 1 m of a unit source. A number of grid cells in the B and C grids emerged as potential sources of instability. These were generally minor indentations of the coastline, barely resolved by the grids, or narrow channels. The region contains several inlets (called Esteros) extending far inland that, at a practical level of spatial resolution, proved difficult to accommodate. Among these are the upper reaches of the multiple arms of Drake's Estero and, feeding into Bodega Bay, Estero Americano and Estero

de San Antonio. A limited number of grid cells in the outermost (A) grid required correction. Generally these were associated with non-physical features in the topographic database, such as where a track of ship-based soundings were improperly merged with other data sources. After an iterative process of grid correction and retesting using these sources, both the reference and forecast model grids were deemed satisfactory and the testing of extreme and historical events could begin. Figure 9 illustrates a step in the process where a deficiency in the reference model grid generated a mild instability (in the EPSZ B19 micro-tsunami scenario –see Table 6) causing the reference model time series at the warning point, initially in close agreement with the forecast model, to develop unrealistic, high frequency oscillations. Though still generally tracking the forecast model result, and not growing without bound, the feature could behave erratically in simulating real events. Modification of the reference model bathymetry eliminated the problem, as seen in the lower panel, and micro-tsunami tests involving other sources (RNSZ B14 and ACSZ B6) did not reveal other issues.

4.2 The "mega-tsunami" tests

The record of tsunami impact on the northern California coast discussed later reveals that sources around the entire periphery of the Pacific can be felt. Indeed the catastrophic Indian Ocean tsunami of 2004 was detectable at Point Reyes, California as it was throughout the global ocean. A broad suite of 19 extreme events (so-called mega-tsunamis) are part of the NCTR protocol for the testing of Pacific basin forecast models, and are described in Table 6. Their locations are included in Figure 10. To simulate each mega-tsunami source, ten A-B pairs of unit sources are used, with an evenly distributed slip of 25 m. As described by Gica et al. (2008), each unit source represents a 100×50 km area of the fault surface with the long axis parallel to the plate boundary. The B-row is shallowest, sloping from a nominal depth of 5 km (unless a depth estimate has been provided by the USGS based on the earthquake catalogs), row-A is deeper, followed by rows Z, Y, X ... where appropriate. Thus, the extreme case sources represent 1000 km long ruptures with a width of 100 km; the corresponding magnitude is Mw 9.3.

Discussion of the entire set in greater detail is provided later in the report, once the validity of the forecast model has been established. Here we focus on a subset of three synthetic cases, highlighted in Figure 10 and Table 6, to contrast the forecast model with the more highly resolved reference model. In Figures 11-13 the reference model results (from the subregion spanned by the forecast model) are shown in the upper panel. The corresponding forecast model results in the lower panel employ the same scale. Insets are used to show the time series (black for reference, red for forecast model) of H, U, and V at the warning point point (the Point Reyes tide gauge). The lagged correlation of H at the reference point is drawn in the lower inset and illustrates that there is generally only a few minutes lag between the time series, with the reference model lagging the forecast model. This behavior is repeated in other scenarios illustrated later in the report. It is a general feature of MOST and is due to the tendency of more finely resolved features in the bathymetry slowing the progress of long waves and arises later in the context of model validation using observations.

The agreement between the reference and forecast model results for the three mega-events is good, both for the maximum amplitude and speed distributions, the reference point time series, and the discrete "snapshots" of the amplitude and vector velocity fields. The earliest waves show the best agreement; later in the solution the reference and forecast model results begin to diverge as multiple reflections with the coastline occur. A qualitative difference between the

solutions is often seen along the straight coastline north of Point Reyes Light (see Figures 1 and 2). The straight shoreline, bounded by rocky headlands at the south and north, supports edge waves that appear most noticeably in the reference model results almost as a standing wave pattern, but generally do not propagate around the headland and into Drake's Bay.

It is noticeable that, in all three of the cases shown, the reference model tends to oscillate longer and have somewhat larger amplitude than does the forecast model though the two solutions are in close agreement for the first few tsunami waves. This is likely a physical reality: the more highly resolved bathymetry and coastline of the reference model providing greater scope for non- linear features or reflected waves to develop as, for example, near the rocky headland west of Bodega Bay. This observation suggests a caveat to operational use of the forecast model: while accurate portrayal of the early history of an event is to be expected, the duration of the event and the amplitude of later waves may be under-estimated. Tide gauge data will be needed to verify this conjecture, which is pursued later in the report.

The snapshot comparisons in the lower panels of Figures 11-12 are quite reasonable, illustrating that the solutions match not just at the reference point. It is worth noting too that, although the ACSZ 56–65 mega-event represents a massive Cascadia tsunami, the scale of the impact to the Point Reyes area (\sim 3m) is not substantially greater than from trans-Pacific locations (KISZ 01–10 off Kamchatka and NTSZ 30–39 near Samoa.) The Crescent City response to the same synthetic Cascadia mega-event exceeds 10m (Arcas and Uslu, 2009). It would appear that the energy propagated along shore to the south, perhaps due to sheltering by Cape Mendocino, is reduced and that perhaps the greatest impact to Marin County may be associated with source regions elsewhere in the Pacific basin.

In Figure 13 the comparison time was intentionally chosen later in the event as a counterexample. While the warning point point amplitudes and the nearby fields of the forecast and reference model may be in reasonable agreement, the broader wave patterns may have substantial phase differences. The comparisons in these lower panels is restricted to the portion of C grid area common to both models, There is a suggestion that the near shore velocity fields at the north and south forecast model boundaries differ somewhat from the reference model for which these are internal points.

Before proceeding to validate the model with historical events, one other synthetic event is usual in the testing protocol: a mild source of magnitude 7.5 at a remote location. A single unit source near Samoa (NTSZ-B36, see Figure 10) is employed and its representation by the reference and forecast model are compared in Figure 14. Such an event results in a response of about 4 cm in Point Reyes sea level and again there is excellent agreement between both model representations in the earlier portion of the event.

The results presented above, for a variety of synthetic events, suggest that the reference and forecast model versions of the model are in good agreement. The match is particularly good in the early stages of a wave train; later, as reflections and other interactions with the coastline occur, the solutions may diverge. The next task is to ascertain whether the models replicate observations from actual tsunami events. Given the manner in which the MOST model is forced, at its boundary (with wave amplitudes and currents not available in real observations), it is not possible to validate the model independently. Rather, as described earlier in Section 2.2, the validation will rely on the results of an external model, based perhaps on DART® observations or on a description of the tsunami source in the literature. As a result, the success of the model in replicating observations within its domain is in part dependent on the adequacy of the forcing employed to represent the actual external wave field. For historical events preceding the

DART® array, the unit source representations are based on seismic observations or coastal tide gauge data. Past experience suggests that, in the far field at least, the propagation solution is not overly sensitive to variation in the unit source weights. Nonetheless, imperfections in forecast model predictions of coastal observations will not necessarily indicate a defect in the model itself. Neither are the tide-gauge observations, available for comparison with model prediction, perfect. They may include noise from wind wave activity, possibly amplified by harbor resonances.

4.3 Model validation: the Honshu-2011 tsunami

In addition to its disastrous impact on the coast of Japan, the Honshu tsunami of March 11, 2011 radiated waves throughout the Pacific basin. Those arriving at nearby DART®s were of unprecedented amplitude and their signal/noise ratios facilitated accurate and early source characterization. Further afield, the waves were detectable at all operational DART®s in the basin and, while major damage was mainly confined to Japan, significant signals were obtained at multiple coastal tide gauges. Prior to this event, the Kuril-2006 tsunami event was the best available for model validation. For the U.S. west coast at least, that role has now been taken by Honshu-2011. The adequacy of the composite propagation solution can be assessed by comparison with the BPR signals from the west coast DART®s. An additional BPR record is available for this purpose: the MARS cabled observatory in Monterey Canyon has, since July 2010, had a pair of bottom pressure sensors at a depth of about 870 m. One is a typical BPR, reporting at the standard DART® 15-sec recording interval. The other is an experimental sensor –the "Nano" (Paros et al., 2011) sampling at 40Hz with enhanced sensitivity. For this report we employ only data from the typical BPR.

The locations of the west coast BPRs, reporting during Honshu-2011, are shown in Figure 15. To the left of the locator chart, the actual and propagation model results, interpolated to the BPR locations, are compared. There is clearly a strong agreement but, even for the earliest waves, there are two points of difference. Firstly the model "waves" (drawn in black) arrive about 8 min early, a difference that is small compared to the transit time of over nine hours. Early arrival in the model is typical and is associated with the limited resolution of the basin-wide bathymetry. Finer-scale features in the actual bathymetry slow down the real wave trains (the red curves). The other feature of the model vs. observation comparison is that the model underestimates the observed signal by about 20% at all locations. In the right-hand panel, the lagged and scaled-up versions of the model time series are seen to be in excellent agreement with observation. Since these results are likely the best obtainable with the current state of the DART® array and inversion methodology, less than perfect agreement between forecast model and observations is not necessarily indicative of a major defect in the forecast model itself.

With that caveat, we proceed with the model validation. The prime location for this purpose is the Point Reyes, California tide gauge itself, the "reference point" for the current model. Other coastal locations termed "warning points" are of interest to the TWCs though they may not have their own dedicated forecast model. In its basic form, SIFT's coastal forecast for warning points is generated by extrapolating offshore values from the propagation solution to the coast using Green's Law. Based on simple assumptions this law indicates that the waves should grow in inverse proportion to the one-fourth power of the depth ratio. The assumption is crude at best and it makes sense that, when a forecast model has been run, the predictions within its domain are likely to be superior to the Green's Law equivalent and should replace them in an

enhanced coastal forecast. For the Point Reyes model, tide gauge observations are available at several points within the domain and, in the case of Honshu-2011, all of these had detectable signals. The auxiliary sites are Bolinas, lying within the C grid but within the lagoon and with only 6-min sampling; San Francisco, Alameda, and Richmond within San Francisco Bay and the model B grid and with 1-min sampling; Arena Cove, near the northern bound of the forecast model A grid and also with 1-min sampling. The results of the comparison may be seen in Figure 16 where the reference (black) and forecast model (red) versions of the model response are compared to the observations (drawn in green.) The model curves have been lagged to facilitate the comparison but, unlike the 20% enhancement needed to bring the propagation results into conformity with the DART® observations (see Figure 15), the amplitude within the forecast model has not been adjusted. For the first six hours of the event, the agreement between observation and model is quite gratifying, particularly at Point Reyes itself and at San Francisco. For Arena Cove the agreement is limited more to the early waves; perhaps as a result of resonance associated with local geometry, the observed response grows and shifts to a higher frequency than appears in the model signals. The reference model solution is a better match in amplitude to the observations from Arena Cove than is the forecast model whose representation of the geometry is quite coarse.

The good agreement between the model amplitude and observation at Point Reyes, particularly for the first wave, is an apparent contradiction of the situation at the offshore sites. There, as noted above in the discussion of Figure 15, the model forcing under-estimated the observations by some 20%. The explanation lies in an unfortunate data loss in the 1-min data stream from the coastal tide gauges during the early part of the event. At most sites the loss was 18 min, 16:03 to 16:20 UTC (slightly longer at San Francisco), as highlighted in yellow in Figure 16. During the event, and in the days that followed, 15-sec data were downloaded from some tide gauges to bridge the gap. At Point Reyes this supplementary 15-sec data was unavailable prior to 18:19 UTC. A few points from the 6-min data stream were employed in the bridging and filtering operations that resulted in the observed Point Reyes time series. Six-minute data badly alias the short times scales of the tsunami waves and, as a result, the apparent close agreement of model amplitude and observation for the first wave peak is probably fortuitous.

Progressing deeper into San Francisco Bay, at Alameda and Richmond, the match between the models themselves and with the observations, is degraded compared with the better agreement near the entrance. Nonetheless the agreement is quite good and shows promise for an improved "coastal forecast" usage of Point Reyes' forecast model results. Least satisfying, but understandable, is the comparison at Bolinas. The tide gauge there lies within the mouth of the lagoon and an adequate representation of the narrow entrance channel is difficult, particularly in the forecast model. As is common with narrow-mouthed entrances to enclosed regions, there is a tendency for the model to retain water within and clearly (the red curve) the lagoon increasingly does not empty during the "ebb" phase of the tsunami wave train. The reference model solution, perhaps as a result of excessive modifications or inaccurate representation of entrance geometry in the DEM, seems to resonate far more than the observational record. It is possible, however that, with its 6-min sampling and placement, the Bolinas tide gauge is not well suited to tsunami detection. On a positive note, the timing and amplitude are not grossly dissimilar to the data. The purpose of the forecast model is more to predict the impact on the seaward side of the Stinson Beach spit and, based on the success at San Francisco, forecasts outside constricted regions of the model domain are likely to be quite useful for warning purposes.

We now step back in order to verify the agreement between the reference and forecast model solutions throughout the common portion of the C grid domain. In Figure 17, as was done for the purely synthetic scenarios, the solutions are compared based on their maximum amplitude and speed fields, and the time series and lagged correlation at the Point Reyes tide gauge site. The distribution patterns of the maximum fields are comparable and, based on the Point Reyes time series sample, it is not unexpected that the reference model should be the greater with the mismatch coming perhaps for the later waves. A point wise (zero lag) correlation distribution (not shown), between reference and forecast model throughout the forecast model C grid domain, indicates that over 60% of the variance is explained except in constricted areas. The lagged correlation inset confirms a phase difference of only a few minutes between the reference and forecast model time series at the tide gauge. As a further means of comparing the reference and forecast model solutions, snapshots of the amplitude and velocity fields are also provided in Figure 17. For both the reference (upper panel) and forecast model (lower panel) a common scale is used. The agreement is particularly close when the comparison time (indicated by the green line) is close to the first peak's arrival at the tide gauge. Two later sample times are shown in Figures 17d,e illustrating that phase differences can increase as the event unfolds.

The analysis of the Honshu-2011 in the Point Reyes model is concluded with an examination of the pattern of inundation in Figure 18. For this purpose the full reference model C grid domain is drawn. The model suggests that, had the waves arrived at or above mean high water (MHW), both the Limantour Spit and much of Stinson Beach and the low-lying parts of Bolinas may have been inundated. In fact, as illustrated in the inset based on the observed water level at the Point Reves tide gauge, the waves barely attained MHW. Though the reporting of the impact on the U.S. may have been somewhat muted, given the gravity of the imagery from Japan, it appears that on the U.S. west coast the main evidence of the tsunami was in excessive currents, notably at Santa Cruz and Crescent City. As designed, with model sea level set at MHW, the forecast erred on the side of conservatism. In the northern portion of the reference model domain (excluded in the forecast model C grid) the greatest response was predicted with inundation of the Doran Beach spit and the Dillon Beach / Lawson's Landing area at the north and south ends of Bodega Bay. Although in reality no actual inundation occurred due to the state of the tide, video clips posted online document strong currents beneath the Lawson's Landing pier, and oscillations of 2-3 ft with 20-min periodicity were reported for Dillon Beach. Examination of the model time series, both the reference and forecast versions, from Bodega Bay (not shown) indicate that the northern and southern portions were rising and falling together, so the large amplitudes responsible for the inundation pattern were not associated with the excitation of an alongshore standing wave mode. Also shown in Figure 18 is the CalEMA Inundation Line, based on an ensemble of synthetic mega-tsunami scenarios. The MHW-based model prediction does impact, albeit at a lesser level, the regions that the CalEMA study identifies as vulnerable.

4.4 Model validation with other preferred historical events

We now proceed to examine how well the reference and forecast model solutions compare with observation for several other historical cases highlighted in Table 1a and Figure 6. These are among the preferred cases in the NCTR protocol to be applied in the validation of Pacific Ocean forecast models. The reference and forecast model time series are inter-compared at

Point Reyes tide gauge, Arena Cove, and San Francisco and validated where possible with observation, and the same representations of maximum amplitude, point wise correlation, and snapshots of the reference and forecast model fields are drawn.

The results, displayed and described below, represent other DART®-detected and well documented recent events: Chile-2010, Samoa-2009, and Kuril-2006, the latter being the first substantial event for which direct observation of the tsunami wave train was available from multiple deep-water DART® sites. These events occurred subsequent to the installation of the tide gauge at Point Reyes. Two pre-DART® cases are included in this section: Alaska-1964 and Unimak-1946 whose large amplitudes caused severe damage to Hawaii and provided the impetus for the establishment of the TWCs. Source characterization for these events is based on the literature, with the source mechanism estimated from the seismic record.

For Chile-2010 the direct comparison of the reference and forecast model results are shown in Figure 19 with satisfactory results both in terms of reference to forecast model intercomparison and agreement with the observed time series at three locations displayed in Figure 20. The amplitude series match well throughout the six-hour period shown and there is strong pointwise correlation throughout the common domain. Comparisons of observations with predictions based on the dedicated forecast models are to be found in the forecast model reports for Arena Cove (Spillane, 2010) and San Francisco (Uslu et al., 2010) and in post-event reports online at the NCTR website. Excellent agreement is seen for Point Reyes and San Francisco, though the leading wave at Point Reyes is over-estimated and the timing of some later features at San Francisco is less than perfect. The observational record at Arena Cove is noisier, though the amplitude of the first wave is captured well by both models.

For Samoa-2009, the equivalent set of results is shown in Figures 21 and 22. Despite the considerably more complex structure of the maximum amplitude field, the forecast model pattern is in good agreement with that from the reference model and the time series for the first few hours agree well. Later the forecast model solution appears to decay faster than that from the reference model. Considering the Point Reyes observations in Figure 22, the reference model is in better agreement with the amplitude of later waves. At San Francisco the situation is less clear, with the reference model perhaps overestimating the observed response, while at Arena Cove neither model (as extracted from the A grid) reproduces the severe ringing evident in the observations.

Figures 23 and 24 represent the Kuril-2006 event. Agreement between the models is strong both for the early and later portions of the record shown. However, in comparisons with observation, the models underestimate later features in the San Francisco observations and, at Arena Cove, the forecast model response decays far too rapidly. At both locations outside the C grid, the predicted maxima are less than 50% of what was observed. To summarize these three events, with weaker impacts than Honshu-2011, the accuracy of a revised "coastal forecast" based on the A and B grids may be reduced. It remains to be demonstrated whether they are significantly better than those based on Green's Law.

The Alaska-1964 and earlier Unimak-1946 events were widely felt along the U.S. west coast, though the greatest impact was to the Hawaiian Islands. The model representations of these major pre-DART® events are illustrated in Figures 25-28. The reference and forecast model representations of the Alaska-1964 event in the C grid are seen (Figure 25) to be in close agreement, both in terms of their maxima and at the arrival of the first wave peak. During the Alaska-1964 event a run-up of 240 cm was reported for Drake's Bay (Point Reyes) with 274 cm at Muir Beach and 113 cm at San Francisco. Arena Cove and Bodega experienced run-ups of 183 and 76 cm

respectively. The maxima of the model time series (Figure 26) are in good agreement with these reports though both the reference and forecast model amplitudes for the first wave at San Francisco are about twice the observed value.

The representation of the Unimak-1946 is also satisfactory with close agreement of the pattern of reference and forecast model maxima, the time series of wave amplitude and velocity at the Point Reyes reference point, and for the instantaneous "snapshots" of these fields an hour after the leading peak (Figure 27). Reported run-ups associated with Unimak-1946 were 240 cm at Arena Cove, 130 cm at Bolinas and 256 cm at Muir Beach; run-up at San Francisco and Alameda were 26 and 20cm respectively. The model hindcasts for San Francisco (Figure 28), are in reasonable agreement with observation, and the model maximum at Point Reyes is consistent with the reported run-up at nearby Bolinas. The model result for Arena Cove considerably under-estimates the reported value. This may be due to the coarse representation of Arena Cove in the A grid of the present model, with the better result for the Alaska-1964 event being fortuitious, or indicative of directionality as a factor in model fidelity. However, for both events, the results support the usefulness of model results beyond the C grid for forecast purposes.

4.5 Other historical simulations of interest at Point Reyes, California

The above analysis has documented good agreement between the forecast model and the slower running reference version. This permits us to simulate the balance of the historical cases in Table 1a, where impacts to the study area have been reported (and the remaining mega-tsunami scenarios) with the forecast model alone. These runs are intended to further validate the stability of the forecast model but also provide some information on the exposure of the region to tsunamis generated at various points on the periphery of the Pacific.

The quality of the modeling of the historical events highlighted above is likely to be the result of good characterizations of the source, based on DART® observations in the case of recent tsunamis or extensive post-event analysis in the case of the historical examples. In the absence of direct and timely observations, the successes of the forecast models are likely to be much reduced. An extreme case in point is the Sanriku-1896 event, a so-called "tsunami-earthquake" (Dudley and Lee, 1998), causing devastating losses in Japan despite its modest magnitude and scant warning in the form of ground motion. A digitized marigram from Sausalito (across the Golden Gate from San Francisco) is available from the WCATWC archives and is drawn in the lower panel of Figure 29. While the timing is reasonably represented, the amplitude considerably underestimates the reported run-up of 10 cm at Sausalito and 20 cm at San Francisco. Reports are not available from nearby locations outside the bay, the closest being 150 cm at Santa Cruz.

For the Andreanof-1957 event reported run-up values of 29 cm at Bodega, 26 cm at San Francisco and 18 cm at Alameda are in reasonable agreement with the model results (35 cm, 46 cm, and 23 cm respectively). No observed time series is available for comparison for this event, nor for East Kuril-1994. For the latter only a 4 cm run-up, reported at Alameda is available for validation; the maximum model amplitude for Alameda at 2.5 cm is in good agreement. For the remaining events in Table 1a, time series are available for more thorough validation and are displayed in Figures 30-36 with limited comment; run-up values from the NGDC database, where available, are added as annotations to the graphics.

The impact of Kamchatka-1952 with Mw 8.6 is available as a marigram from San Francisco (Figure 30). Its amplitude there is well represented by the model, suggesting that a run-up

of 3 m or more may have occurred at Point Reyes and elsewhere in Marin County. For the Chile-1960 Mw 9.3 event, whose San Francisco marigram was also obtained from the WCATWC archive, the character of the observed response is quite different from the model representation. As seen in Figure 31, the model exceeds the observed amplitude response, by a factor of 2–3, and lacks the higher frequency components evident in the observations some hours into the event. The model wave arrived about 20 min early. At Alameda, also within San Francisco Bay, the maximum amplitude of the model, at 68 cm, is about twice the reported run-up of 31 cm. At Stinson Beach the model exaggeration is less severe: 217 cm compared to the observed 152 cm, but is again large (68 cm compared to the observed 25 cm) near Bodega.

Figure 32 presents the validation results for the Andreanof-1996 event. At Point Reyes the agreement is quite good and at Alameda the weak model waves seem to capture some of the features of the observed series. At Arena Cove, however, the signal is far too weak to be visible against the high noise background. For the Peru-2001 (Figure 33) and Hokkaido-2003 (Figure 34) events the validation is quite satisfactory, but for the winter Rat Island-2003 event, as seen in Figure 35, there is considerable noise at the validation sites, limiting the visibility of signals as weak as the model predicts. This event is, however, notable in that, aided by direct observations of bottom pressure from pre-cursors to the DART® array, useful forecasts were provided to inform the Hawaii emergency response. The Tonga-2006 event proved useful for validation of the Point Reyes model, with a strong response, shown in Figure 36, that agrees well with observation.

The year 2007 brought several events with which to validate the model, beginning with the normal thrust earthquake seaward of the Kuril Trench on January 2007. As seen in Figure 37, the model correctly captured the leading trough and amplitude seen at Point Reyes and San Francisco, though at Arena Cove the background noise limits the usefulness of the observations. The Solomon-2007 event hindcast (Figure 38) is reasonably satisfactory, though the signal in both the model and the observations is weak. In August an event off Peru (Figure 39) appears to match well the observations at Point Reyes but, at Arena Cove and San Francisco, while the model seems to capture the amplitude and timing of the early waves, the later portion of the event is less satisfactory. The final event to be treated, among those listed in Table 1a, is the weak winter Chile-2007 event. Not surprisingly, since the forecast amplitudes are very small off California, there is not a lot to be learned from this event displayed in Figure 40.

Several additional events, listed in Table 1b, are available for analysis. Of these, the Sanriku-1896 event has been presented earlier, and the Cape Mendocino tsunami of 1992 as the sole, albeit weak, representative of a Cascadia event, is described in the next Section. The remainder, generally weak in terms of their impact and most occurring in winter where the noise background limits the S / N ratio, are not reported other than to state that all ran without difficulty or evidence of instability.

4.6 The Mendocino earthquake of April 25, 1992

Of special interest to northern California is the Mendocino earthquake of April 25, 1992. This has the distinction of being the most recent substantial thrust event on the Cascadia subduction zone. While strike-slip events are commonplace offshore in this region, as shown in the upper right panel of Figure 41, it is thrust faults that have the potential to generate significant vertical displacements of the sea floor that cause large tsunamis. The epicenter of the 1992 event was on land to the southeast of the plate triple-junction off Cape Mendocino. Uplift on

the order of a meter of a 25 km stretch of the nearshore, between Cape Mendocino and Punta Gorda to the south was evident in a die-off of intertidal organisms, reported by Carver et al. (1994). Presumably extending offshore too, this deformation is not well represented well by either of the southernmost unit sources (ACSZ-A/B65) now available in the propagation database (see Figure 6 where the epicenter is marked by the seismic "beach ball".) A custom source, available from NCTR but not part of the propagation database, is used to model the event for comparison with two digitized marigrams, plotted in the lower panels of Figure 41. These were obtained from the WCATWC archives. The model performs reasonably in representing the leading wave, though the model series had to be delayed by 30 min to achieve alignment. This time offset, greater both in actual time units and as a percentage of travel time, than those typically necessary to adjust trans-basin predictions, may be the result of the coarse representation of the near shore bathymetry. Another possible explanation is that this event, described by González et al. (1995), may have generated a train of coastal-trapped edge waves. Traveling slower than normal tsunami waves taking a deep-water route, the edge waves may have resulted in a delayed arrival and an extended duration for the event. This possibility, and the suggestion that the ACSZ source line ought to be extended at least one unit further south, make this an event worth further study. The reference and forecast models for Point Reyes, and other west coast models (Point Reyes, Eureka, Crescent City, etc.), have a major role in ongoing risk assessment studies for Cascadia.

To summarize the analysis of historical events in the preceding sections, it would appear that the Point Reyes forecast model is capable of producing accurate forecasts for this open coast site on the U.S. west coast. Though the actual waves may be difficult to observe accurately at the tide gauge during winter storms, the objective of producing credible forecasts of sizeable tsunami impacts appears to have been met. Enhanced "coastal forecast" estimates for locations within the Point Reyes B grid can be useful while sites in the A grid (as illustrated by Arena Cove) may have less utility.

4.7 Simulation of the remaining synthetic mega-events

We conclude this section with a summary of other model runs that were made in order to verify its stability, but which provide useful information on the exposure of Point Reyes to potentially hazardous future events within the Pacific. As noted earlier, the sparse instrumental record of actual events needs to be augmented with credible scenarios to permit risk assessment. While not pretending to be a full-blown risk assessment for the Point Reyes and southwest Marin County area, the full set of mega-tsunamis modeled during stability testing can provide some early estimates.

Results for the set of 19 mega-tsunamis, based on the forecast model are presented in Figure 42. Each source is a composite of 20 unit sources from the A and B rows with an evenly distributed slip of 25 m representing an Mw 9.3 event. A color-coded square, drawn at the geometric center of each synthetic source, is used to represent the impact at Point Reyes resulting from that source. The measure of impact employed in Figure 42 is the maximum amplitude of the predicted time series at the reference point. There is not any simple relationship apparent between source orientation, location, or great circle distance to Point Reyes; focusing associated with seafloor features can more than compensate for the decay associated with geometric spreading. It is notable that the greatest impact at Point Reyes comes from trans-basin sources, rather than from those representing Cascadia. The latter apparently beam most of their energy

directly onshore or offshore into the open ocean; arrows normal to the plate boundary are used in Figure 42 as an approximate indicator of main beam direction.

Further results from the suite of mega-event scenarios are presented in Table 7. Seven sites within the C and B grids of the forecast model are represented; the first being Point Reyes tide gauge that was illustrated graphically in Figure 42. Limatour Beach is a well-visited recreational site within the Point Reyes National Seashore; Stinson Beach, adjacent to Bolinas, and Muir Beach are coastal communities between Point Reyes and the southern limit of the forecast model C grid at Point Bonita. Doran Beach and Lawson's Landing represent communities within Bodega Bay, which is only represented in the forecast model B grid. San Francisco, also in the B grid, is included owing to the wealth of tsunami records available there. While Point Reyes has the most instances (10) of having the greatest amplitude among the selected sites, for the mega-events treated here, Muir Beach with 7 and the two overall greatest impacts, is clearly threatened; these results are consistent with the large run-up reported at Muir Beach in the historical record. Lawson's Landing too, with the remaining two cases (one representing the southern end of Cascadia, the other the mid-Aleutians, is clearly at risk and given the inundation that might have resulted had the Honshu-2011 waves arrived under adverse tidal condition (Figure 18) and statements by emergency responders in the "Marin Tsunami" video, Bodega Bay perhaps warrants a dedicated forecast model, though lacking an instrumented reference point. Given the linear geometry and orientation of Bodega and Tomales Bays Version 4 of MOST, which is not limited to north-south and east-west grid lines, should be well suited.FMOST

Finally the set of 19 mega-tsunami scenarios, evaluated here, is an approximate match to the set employed in the CalEMA study that established an inundation line for California. In Figure 43 an ensemble of the inundation predictions by the Point Reyes forecast model is compared with the CalEMA result. The forecast model C grid cells inundated by one or more of the meta-tsunami scenarios are colored red; the CalEMA inundation line is drawn in blue. (The flooded area, in square kilometers, associated with each scenario is included in Table 7.) As underlying topography Figure 43 uses the reference model grid to better indicate coastal indentations. The forecast model provides a reasonable match in most of the threatened areas, particularly the Limantour Spit and Beach areas and Stinson Beach. In some areas, such as Muir Beach, the reduced resolution of the forecast model limits the penetration of flooding there. No attempt has been made to adequately represent Tomales Bay in the forecast model. Its shallowness and the constrictions at its mouth cannot be adequately represented at the spatial resolution necessitated by the run-time constraints for emergency usage.

Conclusions

To conclude, good agreement between observations and model predictions for a subset of historical events, including the recent Honshu-2011 tsunami, has been established and the stability of the model for numerous synthetic events has been demonstrated. In particular the reliability of the forecast model, designed to run rapidly in a real time emergency conditions, has been proven by the favorable comparison with reference model predictions, particularly during the early hours of an event. The model will be included in the SIFT system employed operationally at the Tsunami Warning Centers, and will permit the Point Reyes, California beaches and the communities of Bolinas, Stinson Beach, and Muir Beach to be added to the coastal communities for which forecast capability is available. Additionally it will provide a tool for use in risk assessment studies.

In addition to the scenarios run by the author, and reported here, further tests have been made by other members of the group at NCTR, and will continue to be made by staff at the Warning Centers and others, perhaps in training situations. Among the many related tools developed at NCTR is ComMIT (Titov et al., 2011), the Community Model Interface for Tsunamis, which provides a highly intuitive graphical environment in which to exercise and explore forecast models for any combination of propagation database unit sources. Were any of these avenues to reveal a problem with the model, its origin (most likely in some quirk of the bathymetric files) would be located and corrected then the revised version re-installed for operational use. The development of the forecast system will be a dynamic process, with new models added (and old ones revisited) from the current list of U.S. interests and globally. In the coming years it is expected that further capabilities (for example landslides) will be added as algorithms and methodologies mature.

Acknowledgments

Many members of the NCTR group provided valuable assistance in the production of this report. In particular Diego Arcas edited the first draft for content and style. An anonymous reviewer provided numerous valuable suggestions. Nazila Merati, Yong Wei and Jean Newman performed the SIFT testing reported in Appendix C. CalEMA and other California entities distribute GIS online datasets used in the graphics. The modeling could not proceed without the detailed DEM produced at NGDC by the painstaking combination of numerous bathymetric and topographic surveys. Digitized marigrams for a number of historic events were acquired from the WCATWC archives. This publication is partially funded by the Joint Institute for the Study of the Atmosphere and Ocean (JISAO) under NOAA Cooperative Agreements NA17RJ1232 and NA10OAR4320148. It is JISAO Contribution No. 2088 and PMEL Contribution No. 3401.

Chapter 7

References

- Arcas, D.R. and B. Uslu (2009): A Tsunami Forecast Model for Crescent City, California US Department of Commerce, National Oceanic and Atmospheric Administration, Pacific Marine Environmental Laboratory.
- Barberopoulou, A., J. C. Borrero, B. Uslu, M. R. Legg and C. E. Synolakis (2011): A Second Generation of Tsunami Inundation Maps for the State of California. *Pure and Appl. Geophys.*, 168(11), 2133-2146.
- Carignan, K.S., L.A. Taylor, B.W. Eakins, R.J. Caldwell, D.Z. Friday, P.R. Grothe and E. Lim. (2010): Digital Elevation Models of Central California and San Francisco Bay: Procedures, Data Sources and Analysis. ngdc.noaa.gov/dem/report/download/1220
- Carver, G.A., A.S. Jayko, D.W. Valentine and W.H. Li (1994): Coastal uplift associated with the 1992 Cape Mendocino earthquake, northern California. *Geology*, 22(3), 195-198.
- Dudley, W.C. and Min Lee (1998): *Tsunami!*, University of Hawai'i Press, 362pp.
- Dunbar, P. (2007): Increasing public awareness of natural hazards via the Internet. *Nat. Hazards*, 42(3) doi: 10.1007/s11069-006-9072-3, 529 536.
- Gica, E., M. Spillane, V.V. Titov, C.D. Chamberlin, and J.C. Newman (2008): Development of the forecast propagation database for NOAA's Short-term Inundation Forecast for Tsunamis (SIFT). NOAA Tech. Memo. OAR PMEL-139, NTIS: PB2008-109391, 89 pp.
- González, F.I., K. Satake, E.F. Boss and H.O. Mofjeld (1995): Edge wave and non-trapped modes of the 25 April 1992 Cape Mendocino tsunami. *Pure and Appl. Geophys.*, 144(3-4), 409-426, DOI: 10.1007/BF00874375
- Kanamori, H. and J.J. Cipar (1974): Focal process of the great Chilean earthquake, May 22, 1960. *Phys. Earth Planet. Interiors*, 9(2),128-136.
- Lander, J.F. and P.A. Lockridge (1989): United States tsunamis: (including United States possessions): 1690-1988. US Department of Commerce, NOAA, NESDIS, and NGDC, 1989.
- Loeffler, K. and J. Gesell, editors / cinematographers, (2010): Marin Tsunami: U.S. Geological Survey General Information Product 95 (video). URL: pubs.usgs.gov/gip/95/index.html

- López, A.M., and E.A. Okal (2006): A seismological reassessment of the source of the 1946 Aleutian "tsunami" earthquake. *Geophys. J. Int.*, 165(3), 835âĂŞ849, doi: 10.1111/j.1365-246x.2006.02899.x.
- Niemi, T.M. and N.T. Hall (1996): Historical Changes in the Tidal Marsh of Tomales Bay and Olema Creek, Marin County, California. *J. Coastal Res.*, 12(1): 90-102.
- O'Brien, M.P. (1946): Preliminary Report on Seismic Sea Waves from Aleutian Earthquake of April 1, 1946, *Tech. Rep. HE 116207, Wave Project*, Fluid Mechanics Lab., U. Cal. (Berkeley).
- Paros, J., E. Bernard, J. Delaney, C. Meinig, M. Spillane, P. Migliacio, L. Tang, W. Chadwick, T. Schaad, and S. Stalin (2011): Breakthrough underwater technology holds promise for improved local tsunami warnings. Oceans '11 Paper
- Percival, D.B., D.W. Denbo, M.C. Eble, E. Gica, H.O. Mofjeld, M.C. Spillane, L. Tang, and V.V. Titov (2011): Extraction of tsunami source coefficients via inversion of DART[®] buoy data. *Nat. Hazards*, 58(1), doi: 10.1007/s11069-010-9688-1, 567 590.
- Spillane, M.C., E. Gica, V.V. Titov, and H.O. Mofjeld (2008): Tsunameter network design for the U.S. DART[®] arrays in the Pacific and Atlantic Oceans. NOAA Tech. Memo. OAR PMEL-143, 165 pp.
- Spillane, M.C. (2011): A Tsunami Forecast Model for Arena Cove, California. NOAA OAR Special Report, PMEL Tsunami Forecast Series.
- Tang, L., C. Chamberlin, E. Tolkova, M. Spillane, V.V. Titov, E.N. Bernard, and H.O. Mofjeld (2006): Assessment of potential tsunami impact for Pearl Harbor, Hawaii. NOAA Tech. Memo. OAR pmel-131, ntis: pb2007-100617, 36pp.
- Tang, L., V.V. Titov, and C.D. Chamberlin (2009): Development, testing, and applications of site-specific tsunami inundation models for real-time forecasting. *J. Geophys. Res.*, 114, C12025, doi: 10.1029/2009JC005476.
- Titov, V., and F.I. González (1997): Implementation and testing of the Method of Splitting Tsunami (MOST) model. NOAA Tech. Memo. ERL PMEL-112, NTIS: PB98-122773, NOAA/Pacific Marine Environmental Laboratory, Seattle, WA, 11 pp.
- Titov, V.V., F.I. González, E.N. Bernard, M.C. Eble, H.O. Mofjeld, J.C. Newman, and A.J. Venturato (2005): Real-time tsunami forecasting: Challenges and solutions. *Nat. Hazards*, 35(1) Special Issue, U.S. National Tsunami Hazard Mitigation Program, 41-58.
- Titov, V.V., C. Moore, D.J.M. Greenslade, C. Pattiaratchi, R. Badal, C.E. Synolakis, and U. Kânoğlu (2011): A new tool for inundation modeling: Community Modeling Interface for Tsunamis (ComMIT). *Pure Appl. Geophys.*, 168(11)2121-2131, doi: 10.1007/s00024-011-0292-4.
- Titov, V.V., and C.E. Synolakis (1998). Numerical modeling of tidal wave runup. *J. Waterw. Port Coast. Ocean Eng*, 124(4), 157-171.
- Uslu, B., D. Arcas, V.V. Titov, and A.J. Venturato (2010): A Tsunami Forecast Model for San Francisco, California. NOAA OAR Special Report, PMEL Tsunami Forecast Series: Vol. 3, 88 pp.

Wei, Y., E. Bernard, L. Tang, R. Weiss, V. Titov, C. Moore, M. Spillane, M. Hopkins, and U. Kânoğlu (2008): Real-time experimental forecast of the Peruvian tsunami of August 2007 for U.S. coastlines. *Geophys. Res. Lett.*, 35, L04609, doi: 10.1029/2007GL032250.



FIGURES





Figure 1: The Point Reyes area of west and south Marin County, California.





Figure 2: Extract from the oblique 3-D view of the San Francisco DEM provided by NGDC. The focus is Point Reyes; areas of potential inundation identified by CalEMA are highlighted in red.






Figure 4: Distribution of the historical tsunami sources employed for the development of the Point Reyes forecast model. Those highlighted in red are more extensively investigated using the reference model.




Figure 5: A sample time interval from the Point Reyes tsunami-capable tide gauge, unrelated to tsunami activity. The evolving surface wave spectrum is shown in the lower panel.




Figure 6: The setting of Point Reyes and its nested forecast model grids. The C grids of other west coast forecast models are marked, as are various sites with data available for this study. The closest unit sources of the propagation database lie north of Cape Mendocino, and the epicenter of the most recent Cascadia thrust event is marked.




Figure 7: Nested grid representation for the reference model (RM).




Figure 8: Nested grid representation for the forecast model (FM).




Figure 9: Comparison of the reference (RM) and forecast model (FM) time series at the warning point for three "micro-tsunami" sources in the Western Pacific. The lowest panel illustrates the appearance of model instability before the reference model C grid bathymetry was finalized.




Figure 10: Locations of synthetic tsunami scenarios employed in model development. Three micro-tsunami scenarios, and the magnitude 7.5 case, employ a single unit source; 19 combine ten pairs of unit sources to model mega-tsunamis. Details are provided in Table 6. Cases highlighted in red have both forecast and reference model solutions.




Figure 11: Comparison of reference and forecast model results for the ACSZ 56-65 synthetic mega-tsunami, representing the Cascadia Subduction Zone.

(a) distributions of maximum amplitude in the reference (upper panel) and forecast model (lower panel) results with their time series (reference black, forecast model red) and lagged correlation at the Point Reyes tide gauge as insets.




Figure 11: (b) distributions of maximum speed in the reference (upper panel) and forecast model (lower panel) results with the time series of the vector components at the Point Reyes tide gauge as insets.




Figure 11: (c) comparison of the wave amplitude and currents in the reference (upper panel) and forecast model (lower panel) results at the time indicated in the upper panel inset (the first wave peak).




Figure 11: (d) as in Figure 11c but at the later time when the reference and forecast model solutions have diverged somewhat.




Figure 12: As in Figure 11 but for the synthetic KISZ 01–10 event representing Kamchatka, (a) distributions of maximum amplitude in the reference (upper panel) and forecast model (lower panel) results with their time series (reference black, forecast model red) and lagged correlation at the Point Reyes tide gauge as insets.




Figure 12: (b) distributions of maximum speed in the reference (upper panel) and forecast model (lower panel) results with the time series of the vector components at the Point Reyes tide gauge as insets.




Figure 12: (c) comparison of the wave amplitude and currents in the reference (upper panel) and forecast model (lower panel) results at the time indicated in the upper panel inset (the first wave peak).




Figure 12: (d) as in Figure 12c but for a time near the end of the simulation.




Figure 13: As in Figure 11 but for the synthetic NTSZ 30–39 event representing Samoa, (a) distributions of maximum amplitude in the reference (upper panel) and forecast model (lower panel) results with their time series (reference black, forecast model red) and lagged correlation at the Point Reyes tide gauge as insets.




Figure 13: (b) distributions of maximum speed in the reference (upper panel) and forecast model (lower panel) results with the time series of the vector components at the Point Reyes tide gauge as insets.




Figure 13: (c) comparison of the wave amplitude and currents in the reference (upper panel) and forecast model (lower panel) results at the time indicated in the upper panel inset (the first wave peak).




Figure 13: (d) as in Figure 13c but at the later time when the reference and forecast model solutions have diverged somewhat.




Figure 14: As in Figure 11, but for a synthetic moderate event at NTSZ B36 near Samoa. (a) distributions of maximum amplitude in the reference (upper panel) and forecast model (lower panel) results with their time series (reference black, forecast model red) and lagged correlation at the Point Reyes tide gauge as insets.




Figure 14: (b) distributions of maximum speed in the reference (upper panel) and forecast model (lower panel) results with the time series of the vector components at the Point Reyes tide gauge as insets.




Figure 14: (c) comparison of the wave amplitude and currents in the reference (upper panel) and forecast model (lower panel) results at the time indicated in the upper panel inset (the first major peak).




Figure 14: (d) as in Figure 14c but at the later time when the reference and forecast model solutions have diverged somewhat.




Figure 15: Observed time series (red curves) from DART[®] and MARS bottom pressure sensors during the Honshu-2011 event, compared with the forecast model representation (black curves) based on the propagation database (see Table 1((a). Model time series in the right hand panel have been lagged, and a common scale factor of 1.2 applied.




Figure 16: Comparison of observations with reference and forecast model-predicted time series at selected locations where tide gauge data are available:

(a) Point Reyes, Arena Cove, and San Francisco $_55$ The time period highlighted in yellow marks an outage of 18 min or more that occured for the 1-min data streams along the entire west coast during the event. Runup values from the NGDC catalog, when available, are indicated in the lower right of each panel for this and subsequent figures.




Figure 16: (b) Bolinas (6-min data), Alameda, and Richmond.




Figure 17: Intercomparison of reference (upper panel) and forecast model (lower panel) predictions for the Honshu-2011 event.

(a) maximum amplitude and reference point time series in the style of Figure 11a,




Figure 17: (b) maximum speed and reference point current time series in the style of Figure 11b




Figure 17: (c) snapshot of amplitude and velocity at the leading wave peak in the style of Figure 11c,




Figure 17: (d) snapshot of amplitude and velocity at a later wave peak.




Figure 17: (e) snapshot of amplitude and velocity at a later wave trough.




Figure 18: Inundation forecast from the reference model (RM) C grid for the Honshu-2011 event, compared with the CalEMA inundation line. The inset in the upper right shows tide gauge data from Point Reyes. Actual tides were well below MHW so the inundation forecast was overly conservative.




Figure 19: As in Figure 17 but for the Chile-2010 historical event, (a) maximum amplitude and reference point time series in the style of Figure 11a,




Figure 19: (b) maximum speed and reference point current time series in the style of Figure 11b




Figure 19: (c) snapshot of amplitude and velocity at the leading wave peak in the style of Figure 11c.




Figure 20: Model and observed time series comparison for the Chile-2010 event.




Figure 21: As in Figure 17 but for the Samoa-2009 historical event, (a) maximum amplitude and reference point time series,




Figure 21: (b) maximum speed and reference point current time series,




Figure 21: (c) snapshot of amplitude and velocity at a time between waves at the reference point.




Figure 22: Model and observed time series comparison for the Samoa-2009 event.




Figure 23: As in Figure 17 but for the Kuril-2006 historical event, (a) maximum amplitude and reference point time series,




Figure 23: (b) maximum speed and reference point current time series,




Figure 23: (c) snapshot of amplitude and velocity at the leading wave peak.




Figure 24: Model and observed time series comparison for the Kuril-2006 event.




Figure 25: As in Figure 17 but for the Alaska-1964 historical event, (a) maximum amplitude and reference point time series,




Figure 25: (b) maximum speed and reference point current time series,




Figure 25: (c) snapshot of amplitude and velocity at the leading wave peak.




Figure 26: Model and observed time series comparison for the Alaska-1964 event.




Figure 27: As in Figure 17 but for the Unimak-1946 historical event, (a) maximum amplitude and reference point time series,




Figure 27: (b) maximum speed and reference point current time series,




Figure 27: (c) snapshot of amplitude and velocity at the later wave peak.




Figure 28: Model and observed time series comparison for the Unimak-1946 event.




Figure 29: The Sanriku event of June 15, 1896.




Figure 30: The Kamchatka event of November 4, 1952.




Figure 31: The Chile event of May 22, 1960.




Figure 32: The Andrean of event of June 10, 1996. $\,\,86$




Figure 33: The Peru event of June 23, 2001.




Figure 34: The Hokkaido event of September 25, 2003.




Figure 35: The Rat Island event of November 17, 2003.




Figure 36: The Tonga event of May 3, 2006.




Figure 37: The normal thrust event off the Kuril Islands on January 13, 2007.




Figure 38: The Solomon event of April 1, 2007.




Figure 39: The Peru eyent of August 15, 2007.




Figure 40: The Chile event of November 14, 2007.




Figure 41: The Cape Mendocino event of April 25, 1992. The upper panels show the frequency of non-thrust events in the vicinity, with only two having a focal mechanism characteristic of subduction. Lower panel: comparison of model with observation at Arena Cove and Point Reyes.




Figure 42: Predicted maximum sea level (from the forecast model) at the Point Reyes tide gauge that might result were "mega-tsunamis" to occur at various locations around the Pacific basin.




Figure 43: Chart of the area inundated by one or more of the mega-tsunami scenarios based on the forecast model. Shown in blue is the CalEMA inundation line, which is based on a similar ensemble of scenarios.

TABLES



	Earthquake / Seismic	Seismic				Model
	USGS	CMT				
Event	Date Time (UTC) Epicenter	Date Time (UTC) Centroid	Magnitude Mw	Tsunami Magnitude	Subduction Zone	Tsunami Source (Reference/Derivation)
1946 Unimak	01 Apr 12:28:56 52.75°N 163.50°W	Not Available	8.5	8.5	ACSZ	7.5×B23 + 19.7×B24 + 3.7×B25 (López & Okal, 2006)
1952 Kamchatka	04 Nov 16:58:26.0 52.76°N 160.06°E	Not Available	9.0	0.6	KISZ	$19.71 \times (A4 + Y4 + Z4 + A5 + Y5 + Z5 + A6 + Y6 + Z6)$ (ad hoc)
1957 Andreanof	09 Mar 14:22:31 51.56°N 175.39°W	Not Available	8.6	8.7	ACSZ	31.4×A15 + 10.6×A16 + 12.2×A17 (Preliminary)
1960 Chile	22 May 19:11:14 38.29°S 73.05°W	Not Available	9.5	9.5	CSSZ	125×(A93 + B93 + Z93 + A94 + B94 + Z94 + A95 + B95) Kanamori & Cipar (1974)
1964 Alaska	28 Mar 03:36:00 61.02°N 147.65°W	Not Available	9.2	6.8	ACSZ	15.4xA34 + 18.3xB34 + 48.3xZ34 +19.4xA35 +15.1xB35 (Tang et al., 2006, 2009)
1994 East Kuril	04 Oct 13:22:58 43.73°N 147.321°E	04 Oct 13:23:28.5 43.60°N 147.63°E	8.3	8.1	KISZ	9.0×A20 (ad hoc)
1996 Andreanof	10 Jun 04:03:35 51.56°N 175.39°W	10 Jun 04:04:03.4 51.10°N 177.410°W	7.9	7.8	ACSZ	2.40×A15 + 0.80×B16 (Preliminary)
2001 Peru	23 Jun 20:33:14 16.265°S 73.641°W	23 Jun 20:34:23.3 17.28°S 72.71°W	8.4	8.2	CSSZ	5.7×A15+2.9×B16+1.98×A16 (Preliminary)
2003 Hokkaido	25 Sep 19:50:06 41.775°N 143.904°E	25 Sep 19:50:38.2 42.21°N 143.84°E	8.3	8.3	KISZ	$3.95 \times (A22 + B22 + A23 + B23)$ (ad hoc)
2003 Rat Island	17 Nov 06:43:07 51.13°N 178.74°E	17 Nov 06:43:31.0 51.14°N 177.86°E	7.7	7.8	ACSZ	2.81×B11 (Real-time)
2006 Tonga	03 May 15:26:39 20.13°S 174.161°W	03 May 15:27:03.7 20.39°S 173.47°W	8.0	8.0	NTSZ	6.6×B29 (ad hoc)
2006 Kuril	15 Nov 11:14:16 46.607°N 153.230°E	15 Nov 11:15:08 46.71°N 154.33°E	8.3	8.1	KISZ	4.0xA12+0.5xB12+2.0xA13+1.5xB13 (Real-time)
2007 Kuril	13 Jan 04:23:20 46.272°N 154.455°E	13 Jan 04:23:48.1 46.17°N 154.80°E	8.1	7.9	KISZ	-3.64 × B13 (Real-time)
2007 Solomon	01 Apr 20:39:56 8.481°S 156.978°E	01 Apr 20:40:38.9 7.76°S 156.34°E	8.1	8.2	NVSZ	12.0×B10 (Preliminary)
2007 Peru	15 Aug 23:40:57 13.354°S 76.509°W	15 Aug 23:41:57.9 13.73°S 77.04°W	8.0	8.1	CSSZ	$0.9 \times A61 + 1.25 \times B61 + 5.6 \times A62 + 6.97 \times B62 + 3.5 \times Z62$ (Preliminary)
2007 Chile	14 Nov 15:40:50 22.204°S 69.869°W	14 Nov 15:41:11.2 22.64°S 70.62°W	7.7	7.6	CSSZ	1.65×Z73 (Real-time)
2009 Samoa	29 Sep 17:48:10 15.509°S 172.034°W	29 Sep 17:48:26.8 15.13°S 171.97°W	8.1	8.1	NTSZ	3.96×A34 + 3.96×B34 (Real-time)
2010 Chile	27 Feb 06:34:14 35.909°S 72.733°W	27 Feb 06:35:15.4 35.95°S 73.15°W	8.8	8.8	CSSZ	17.24×A88 + 8.82×A90 + 11.84×B88 + 18.39×B89 + 16.75×B90 + 20.78×Z88 + 7.06×Z90 (Real-time)
2011 Honshu	11 Mar 05:46:24 38.297°N 142.372°E	11 Mar 05:47:47.2 38.486°N 142.597°E	9.0	0.6	KISZ	4.66 × B24 + 12.23 × B25 + 26.31 × A26 + 21.27 × B26 + 22.75 × A27 + 4.98 × B27 (Real-time; Tang et al., 2012)

Table 1: Source characterization for historical tsunami events employed in Point Reyes model testing. Those in bold text were used to compare the reference and forecast model versions. Sources identified as "preliminary" or "ad hoc" may not be identically defined in other forecast model reports.

a) The standard set for Pacific Ocean models.

	Earthquake / Seismic	e / Seismic			Tsunami Source (R	Tsunami Source (Reference/Derivation)
	nses	CMT				
Event	Date Time (UTC) Epicenter	Date Time (UTC) Centroid Magnitude	Magnitude M.,	Tsunami Magnitude	Subduction Zone	Tsunami Source
1896 Sanriku	15 Jun 10:33:00 39.5°N		7.6	7.6	KISZ	b25 x 1.413
	144.0°E					(ad hoc)
1992	25 Apr 18:06:04 40.368°N	25 Apr 18:06:11.8 38.56°N	7.2	7.2	ACSZ	a65 x 0.355 or b65 x 0.355
Mendocino	124.316°W	123.31°W				(ad hoc)
1995 Chile	30 Jul 05:11:24 23.340°S	30 Jul 05:11:56,9 24,17°S	8.0	8.0	CSSZ	$2.812 \times (a75 + b75)$
	70.294°W	70.74°W				(ad hoc)
1995 Kuril	03 Dec 18:01:09 44.663°N	03 Dec 18:01:36.1 44.82°N	7.9	7.9	KISZ	$1.991 \times (a17 + z17)$
	149.300°E	150.17°E				(ad hoc)
1996 Irian	17 Feb 05:59:31 0.891°S	17 Feb 06:00:02.8 0.67°S	8.2	8.2	NGSZ	2.7984 x (a9 + b9 + a10 + b10)
Jaya	136.952°E	136.62°E				(ad hoc)
2009 PapuaNG	03 Jan 19:43:51 0.414°S	03 Jan 19:44:09.0 0.38°S	7.6	7.6	NGSZ	$0.7046 \times (b13 + b14)$
	132.885°E	132.83°E				(ad hoc)
2009 Kuril	15 Jan 17:49:39 46.857°N	15 Jan 17:49:48.3 46.97°N	7.4	7.4	KISZ	b12 x 0.7063
	155.154°E	155,39°E				(ad hoc)
2009 Vanuatu	07 Oct 22:03:15 13:052°S	07 Oct 22:03:28.9 12.59°S	7.6	7.6	NVSZ	1.2×B24 + 0.26×A23
,	166.187°E	166.27°E				followed after 15minutes by
Santa Cruz	07 Oct 22:18:26 12:554°S	07 Oct 22:19:15.3 11.86°S	7.8	7.9	NVSZ	2.6xB23 + 0.9xA23
	166.320°E	166.01°E				(Preliminary; Yong Wei, Personal
						Comministion

Table 1: b) Supplementary historical tsunami events employed for Point Reyes, California forecast model testing.

Grid Area	San Francisco, California
Coverage Area	123.30° to 121.85° W; 37.32° to 38.48° N
Coordinate System	Geographic decimal degrees
Horizontal Datum	World Geodetic System 1984 (WGS84)
Vertical Datum	Mean High Water (MHW)
Vertical Units	Meters
Cell Size	1/3 arc-second
Grid Format	ESRI Arc ASCII grid
Version Employed	Update of February 24, 2011

Table 2: The main features of the San Francisco digital elevation model (DEM), which includes Point Reyes, California.

Point Reyes, California	Sta	ntion#9415020	37°59.7'N, 122°58.6'W			
Tidal Da	tum and Ra	nge Values (Epoch 198	33-2001)			
MHHW (Mean Higher High)	2.964m					
MHW (Mean High Water)	2.760m					
MSL (Mean Sea Level)	2.152m	Great Diurnal Range	Mean Range 1.193m			
MLW (Mean Low Water)	1.567m	1.758m				
MLLW (Mean Lower Low)	1.206m					
	Sea Level Trends and Cycles					
Long Term SL Trend	Increasing 2.10±1.52mm/year					
Seasonal Cycle Range	Minim	um -89mm(April); Max	kimum 59mm(September)			
Interannual Variation	Min	imum -20mm(1989); M	faximum +21mm(1998)			
(from1980)						
	Extremes	to date (June 2011)				
Maximum		3.810m on 06	Feb, 1998			
Minimum		0.387 on 19 J	Jan, 1988			



Table 3: Tidal characteristics of the Point Reyes, California tide gauge (9415020).

Point Reyes, California Reference Model (RM)

Minimum offshore depth 1.5 m; Water depth for dry land 0.1m; Friction Coefficient (n^2) 0.0009 CPU time for a 4-hr simulation 2.69 hr

(Grid	Zonal	Extent	Meridion	al Extent	Resolution	Grid Points
	A	128.000°W	121.500°W	36.000°N	42.500°N	30"x30"	781 x 781
	В	123.300°W	122.100°W	37.475°N	38.475°N	4"x3"	1081 x 1201
	C	123.150°W	122.533°W	37.825°N	38.350°N	4/3"x1"	1666 x 1891

Point Reyes, California Forecast Model (FM)

Minimum offshore depth 2.5 m; Water depth for dry land 0.1m; Friction Coefficient (n²) 0.0009 CPU time for a 4-hr simulation 8 min

		CI C tim	c for a + in 5	illialation o il	1111	
Grid	Zonal	Extent	Meridion	al Extent	Resolution	Grid Points
A	125.000°W	122.000°W	37.00°N	39.00°N	60"x60"	181 x 121
В	123.300°W	122.100°W	37.550°N	38.475°N	18" x 15"	241 x 233
C	123.130°W	122.533°W	37.825°N	38.100°N	4" x 3"	538 x 331



Table 4: Specifics of the reference (RM) and forecast model (FM) grids employed for Point Reyes, California. For the paired values in the resolution and grid points columns, the zonal (East to West) value is listed first, followed by meridional (North to South). CPU times for a 4-hr simulation are based on use of a single Intel[®] Xeon[®] E5670 2.93GHz processor.

Grid	Filename	Maximum	Minimum	Model Time	Water
Ond	Thename	Depth (m)	CFL (s)	Step (s)	Cells
Α	PtReyesCA_RM_A	5002	3.350	1.2 (2x)	436,723
Λ	PtReyesCA_FM_A	4379	7.137	6.0 (3x)	15,977
В	PtReyesCA_RM_B	2166	0.637	0.6 (1x)	664,682
В	PtReyesCA_FM_B	2114	3.062	2.0 (1x)	26,598
C	PtReyesCA_RM_C	98.6	0.995	0.6	1,411,698
	PtReyesCA_FM_C	94.7	3.045	2.0	103,086

Table 5: Grid file names and grid-related parameters. The time steps for the A and B grids must be integer multiples of the basic time step chosen for the C grid.

Scenario Name	Source Zone	Tsunami Source	α [m]
Hicro-tsunami Scenarios EPSZ B19 East Philippines B19 RNSZ B14 Ryukus-Kyushu-Nankai B14 ACSZ B6 Aleutian-Alaska-Cascadia B6 Mega-tsunami Scenarios KISZ 1-10 Kamchatka-Yap-Mariana-Izu-Bonin A2-A31, B22-B31 KISZ 32-41 Kamchatka-Yap-Mariana-Izu-Bonin A32-A41, B32-B41 KISZ 56-65 Kamchatka-Yap-Mariana-Izu-Bonin A56-A65, B56-B65 ACSZ 6-15 Aleutian-Alaska-Cascadia A6-A15, B6-B15 ACSZ 16-25 Aleutian-Alaska-Cascadia A16-A25, B16-B25 ACSZ 22-31 Aleutian-Alaska-Cascadia A22-A31, B22-B31 ACSZ 50-59 Aleutian-Alaska-Cascadia A50-A59, B50-B59 ACSZ 56-65 Aleutian-Alaska-Cascadia A50-A59, B50-B59 ACSZ 56-65 Aleutian-Alaska-Cascadia A56-A65, B56-B65 CSSZ 1-10 Central and South America A1-A10, B1-B10 CSSZ 37-46 Central and South America A37-A46, B37-B46 CSSZ 89-98 Central and South America A37-A46, B37-B46 CSSZ 89-98 Central and South America A89-B98, B89-B98 CSSZ 102-111 Central and South America A102-A111, B102- NTSZ 30-39 New Zealand-Kermadec-Tonga A30-A39, B30-B39 NVSZ 28-37 New Britain-Solomons-Vanuatu A28-A37, B28-B37 MOSZ 1-10 ManusOCB A1-A10, B1-B10 NGSZ 3-12 North New Guinea A3-A12, B3-B12 EPSZ 6-15 East Philippines A6-A15, B6-B15 RNSZ 12-21 Ryukus-Kyushu-Nankai A12-A21, B12-B21 Mw 7.5 Scenario NTSZ B36 New Zealand-Kermadec-Tonga B36			
EPSZ B19	East Philippines	B19	0.01
RNSZ B14	Ryukus-Kyushu-Nankai	B14	0.01
ACSZ B6	Aleutian-Alaska-Cascadia	В6	0.01
	Mega-tsunami Scena	arios	
KISZ 1-10	Kamchatka-Yap-Mariana-Izu-Bonin	A1-A10, B1-B10	25
KISZ 22-31	Kamchatka-Yap-Mariana-Izu-Bonin	A22-A31, B22-B31	25
KISZ 32-41	Kamchatka-Yap-Mariana-Izu-Bonin	A32-A41, B32-B41	25
KISZ 56-65	Kamchatka-Yap-Mariana-Izu-Bonin	A56-A65, B56-B65	25
ACSZ 6-15	Aleutian-Alaska-Cascadia	A6-A15, B6-B15	25
ACSZ 16-25	Aleutian-Alaska-Cascadia	A16-A25, B16-B25	25
ACSZ 22-31	Aleutian-Alaska-Cascadia	A22-A31, B22-B31	25
ACSZ 50-59	Aleutian-Alaska-Cascadia	A50-A59, B50-B59	25
ACSZ 56-65	Aleutian-Alaska-Cascadia	A56-A65, B56-B65	25
CSSZ 1-10	Central and South America	A1-A10, B1-B10	25
CSSZ 37-46	Central and South America	A37-A46, B37-B46	25
CSSZ 89-98	Central and South America	A89-B98, B89-B98	25
CSSZ 102-111	Central and South America	A102-A111, B102-	25
NTSZ 30-39	New Zealand-Kermadec-Tonga	A30-A39, B30-B39	25
NVSZ 28-37	New Britain-Solomons-Vanuatu	A28-A37, B28-B37	25
MOSZ 1-10	ManusOCB	A1-A10, B1-B10	25
NGSZ 3-12	North New Guinea	A3-A12, B3-B12	25
EPSZ 6-15	East Philippines	A6-A15, B6-B15	25
RNSZ 12-21	Ryukus-Kyushu-Nankai	A12-A21, B12-B21	25
	M _W 7.5 Scenario	1	
NTSZ B36	New Zealand-Kermadec-Tonga	B36	1

Table 6: Synthetic tsunami events employed in Point Reyes, California model testing. The reference and forecast model solutions of those shown in bold text were inter-compared extensively.

Scenario Name	Floo	ding				npact S				Amp.
(Great Circle, km)	Area	Rank	PTR ¹	LIM ²	STN ³	MUR ⁴	DOR ⁵	LAW ⁶	SFO ⁷	Rank
AC 56-65 (688)	5.18	7	159	152	160	182	201	224	115	13
AC 50-59 (1278)	4.72	11	202	106	217	373	194	193	203	4
CS 1-10 (2994)	1.18	18	99	69	64	72	48	52	37	18
AC 22-31 (3277)	6.34	4	239	221	288	227	251	333	150	6
AC 16-25 (3731)	4.97	8	266	121	234	275	162	194	102	8
AC 6-15 (4731)	2.55	17	134	87	117	136	118	120	81	17
KI 1-10 (5856)	4.93	10	354	152	184	245	144	189	90	6
CS 37-46 (6070)	0	19	42	36	38	35	37	38	25	19
KI 22-31 (7724)	4.24	12	251	129	170	231	212	182	74	11
NT 30-39 (8054)	7.00	2	402	226	263	277	239	309	127	3
KI 32-41 (8368)	6.39	3	318	169	288	502	361	440	159	2
RN 12-21 (8808)	3.27	15	209	84	115	162	110	121	57	14
KI 56-65 (9429)	3.94	13	166	96	145	233	171	204	87	12
NV 28-37 (9553)	4.96	9	258	131	149	149	173	202	88	10
MO 1-10 (9943)	7.71	1	460	295	324	513	240	277	200	1
CS 89-98 (10063)	3.48	14	140	134	102	78	102	136	43	16
NG 3-12 (10801)	3.15	16	162	107	143	145	133	131	104	15
EP 6-15 (10932)	6.31	5	246	160	264	296	211	235	137	7
CS 102-111 (11010)	3.27	6	265	132	156	193	157	172	77	9
	Overa	ll Max	402	295	324	513	361	440	203	

C-Grid: 1-Point ReyesTG, 2-Limantour Beach, 3-Stinson Beach, 4-Muir Beach, B-Grid: 5-Doran Spit, 6-Lawson's Landing, 7-San Francisco TG.

Table 7: Mega-tsunami scenario impacts, represented by flooding (km^2) in the sub-region $123.1^{\circ}-122.5^{\circ}$ W, $37.8^{\circ}-38.1^{\circ}$ N) and maximum amplitude (cm) at several sites within the model domain (identified in the footnote). The maxima are highlighted and ranked.

Appendix A

Model input files for Point Reyes, California.

As discussed in Section 3.5, input files providing model parameters, the file names of the nested grids, and the output specifications are necessary in order to run the model in either its reference or forecast mode. These files are provided below; each record contains the value(s) and an annotation of purpose.



A.1 Reference model *.in file for Point Reyes, California

The following table contains the parameter and file choices used in the input file for the SIFT implementation (most3_facts_nc.in) of the reference model (RM) for Point Reyes, California. When run on an Intel[®] Xeon[®] E5670 2.93GHz processor during development the model simulated four hours in 2.69 CPU hours.

0.001	Minimum amp. of input offshore wave (m)
1.5	Minimum depth of offshore (m)
0.1	Dry land depth of inundation (m)
0.0009	Friction coefficient (n**2)
1	Let A-Grid and B-Grid run up
900.0	Max eta before blow-up (m)
1.0	Time step (sec)
28800	Total number of time steps in run
2	Time steps between A-Grid computations
1	Time steps between B-Grid computations
30	Time steps between output steps
0	Time steps before saving first output step
1	Save output every n-th grid point
PtReyesCA_RM_A.most	A-grid bathymetry file
PtReyesCA_RM_B.most	B-grid bathymetry file
PtReyesCA_RM_C.most	C-grid bathymetry file
./	Directory of source files
./	Directory for output files
1111	netCDF output for A, B, C, SIFT
1	Number of time series locations
3 335 967	Grid & cell indices for Reference Point

A.2 Forecast model *.in file for Point Reyes, California

The following table contains the parameter and file choices used in the input file for the SIFT implementation (most3_facts_nc.in) of the optimized forecast model for Point Reyes, California. When run on an Intel[®] Xeon[®] E5670 2.93GHz processor the model simulates four hours in under 8 min, satisfying the 10 min target for this metric.

0.001	Minimum amp. of input offshore wave (m)
2.5	Minimum depth of offshore (m)
0.1	Dry land depth of inundation (m)
0.0009	Friction coefficient (n**2)
1	Let A-Grid and B-Grid run up
900.0	Max eta before blow-up (m)
2.0	Time step (sec)
32400	Total number of time steps in run
3	Time steps between A-Grid computations
1	Time steps between B-Grid computations
15	Time steps between output steps
0	Time steps before saving first output step
1	Save output every n-th grid point
PtReyesCA_FM_A.most	A-grid bathymetry file
PtReyesCA_FM_B.most	B-grid bathymetry file
PtReyesCA_FM_C.most	C-grid bathymetry file
./	Directory of source files
./	Directory for output files
1111	netCDF output for A, B, C, SIFT
1	Number of time series locations
3 139 125	Grid & cell indices for 237.02333333 37.99666667

Appendix B

Propagation Database: Pacific Ocean Unit Sources

NOAA Propagation Database presented in this section is the representation of the database as of March, 2013. This database may have been updated since March, 2013.




Figure B.1: Aleutian–Alaska–Cascadia Subduction Zone unit sources.

 ${\bf Table~B.1:} \ Earth quake~parameters~for~Aleutian-Alaska-Cascadia~Subduction~Zone~unit~sources.$

Segment	Description	Longitude(^o E)	Latitude(^o N)	Strike(0)	Dip(o)	Depth (kn
acsz-1a	Aleutian-Alaska-Cascadia	164.7994	55.9606	299	17	19.61
acsz–1b	Aleutian–Alaska–Cascadia	164.4310	55.5849	299	17	5
acsz–2a	Aleutian–Alaska–Cascadia	166.3418	55.4016	310.2	17	19.61
acsz–2b	Aleutian–Alaska–Cascadia	165.8578	55.0734	310.2	17	5
acsz–3a	Aleutian–Alaska–Cascadia	167.2939	54.8919	300.2	23.36	24.82
acsz–3b	Aleutian–Alaska–Cascadia	166.9362	54.5356	300.2	23.36	5
acsz–4a	Aleutian–Alaska–Cascadia	168.7131	54.2852	310.2	38.51	25.33
acsz–4b	Aleutian–Alaska–Cascadia	168.3269	54.0168	310.2	24	5
acsz–5a	Aleutian–Alaska–Cascadia	169.7447	53.7808	302.8	37.02	23.54
acsz–5b	Aleutian–Alaska–Cascadia	169.4185	53.4793	302.8	21.77	5
acsz–6a	Aleutian–Alaska–Cascadia	171.0144	53.3054	303.2	35.31	22.92
acsz–6b	Aleutian–Alaska–Cascadia	170.6813	52.9986	303.2	21	5
acsz–7a	Aleutian–Alaska–Cascadia	172.1500	52.8528	298.2	35.56	20.16
acsz–7b	Aleutian–Alaska–Cascadia	171.8665	52.5307	298.2	17.65	5
acsz–8a	Aleutian–Alaska–Cascadia	173.2726	52.4579	290.8	37.92	20.35
acsz–8b	Aleutian–Alaska–Cascadia	173.0681	52.1266	290.8	17.88	5
acsz–9a	Aleutian–Alaska–Cascadia	174.5866	52.1434	289	39.09	21.05
acsz–9b	Aleutian–Alaska–Cascadia	174.4027	51.8138	289	18.73	5
acsz–10a	Aleutian–Alaska–Cascadia	175.8784	51.8526	286.1	40.51	20.87
acsz–10b	Aleutian–Alaska–Cascadia	175.7265	51.5245	286.1	18.51	5
acsz–11a	Aleutian–Alaska–Cascadia	177.1140	51.6488	280	15	17.94
acsz–11b	Aleutian–Alaska–Cascadia	176.9937	51.2215	280	15	5
acsz–12a	Aleutian–Alaska–Cascadia	178.4500	51.5690	273	15	17.94
acsz–12b	Aleutian–Alaska–Cascadia	178.4130	51.1200	273	15	5
acsz-13a	Aleutian-Alaska-Cascadia	179.8550	51.5340	271	15	17.94
acsz-13b	Aleutian-Alaska-Cascadia	179.8420	51.0850	271	15	5
acsz–14a	Aleutian-Alaska-Cascadia	181.2340	51.5780	267	15	17.94
acsz–14b	Aleutian-Alaska-Cascadia	181.2720	51.1290	267	15	5
acsz-15a	Aleutian-Alaska-Cascadia	182.6380	51.6470	265	15	17.94
acsz-15b	Aleutian-Alaska-Cascadia	182.7000	51.2000	265	15	5
acsz-16a	Aleutian-Alaska-Cascadia	184.0550	51.7250	264	15	17.94
acsz-16b	Aleutian-Alaska-Cascadia	184.1280	51.2780	264	15	5
acsz–17a	Aleutian-Alaska-Cascadia	185.4560	51.8170	262	15	17.94
acsz–17b	Aleutian-Alaska-Cascadia	185.5560	51.3720	262	15	5
acsz-18a	Aleutian-Alaska-Cascadia	186.8680	51.9410	261	15	17.94
acsz-18b	Aleutian-Alaska-Cascadia	186.9810	51.4970	261	15	5
acsz–19a	Aleutian-Alaska-Cascadia	188.2430	52.1280	257	15	17.94
acsz–19b	Aleutian-Alaska-Cascadia	188.4060	51.6900	257	15	5
acsz–20a	Aleutian–Alaska–Cascadia	189.5810	52.3550	251	15	17.94
acsz-20b	Aleutian-Alaska-Cascadia	189.8180	51.9300	251	15	5
acsz–21a	Aleutian-Alaska-Cascadia	190.9570	52.6470	251	15	17.94
acsz-21b	Aleutian-Alaska-Cascadia	191.1960	52.2220	251	15	5
acsz–21z	Aleutian-Alaska-Cascadia	190.7399	53.0443	250.8	15	30.88
acsz–22a	Aleutian–Alaska–Cascadia	192.2940	52.9430	247	15	17.94
acsz–22b	Aleutian–Alaska–Cascadia	192.5820	52.5300	247	15	5
acsz–22z	Aleutian-Alaska-Cascadia	192.0074	53.3347	247.8	15	30.88
acsz-23a	Aleutian-Alaska-Cascadia	193.6270	53.3070	245	15	17.94
acsz-23b	Aleutian-Alaska-Cascadia	193.9410	52.9000	245	15	5
acsz–23z	Aleutian-Alaska-Cascadia	193.2991	53.6768	244.6	15	30.88
acsz–24a	Aleutian-Alaska-Cascadia	194.9740	53.6870	245	15	17.94
acsz–24b	Aleutian-Alaska-Cascadia	195.2910	53.2800	245	15	5
acsz-24y	Aleutian-Alaska-Cascadia	194.3645	54.4604	244.4	15	43.82
acsz–24z	Aleutian-Alaska-Cascadia	194.6793	54.0674	244.6	15	30.88
acsz–25a	Aleutian–Alaska–Cascadia	196.4340	54.0760	250	15	17.94
acsz–25b	Aleutian–Alaska–Cascadia	196.6930	53.6543	250	15	5
acsz–25y	Aleutian–Alaska–Cascadia	195.9009	54.8572	247.9	15	43.82
acsz–25z	Aleutian–Alaska–Cascadia	196.1761	54.4536	248.1	15	30.88
acsz-26a	Aleutian–Alaska–Cascadia	197.8970	54.3600	253	15	17.94
acsz–26b	Aleutian–Alaska–Cascadia	198.1200	53.9300	253	15	5
acsz-26y	Aleutian–Alaska–Cascadia	197.5498	55.1934	253.1	15	43.82
400L LUY	maska Gascadia	101.0100	00.1001	200.1	10	10.02

Table B.1 – continued

		Table B.1 – cor	itinuea			
Segment	Description	Longitude(°E)	Latitude(^o N)	Strike(0)	Dip(o)	Depth (km
acsz-26z	Aleutian-Alaska-Cascadia	197.7620	54.7770	253.3	15	30.88
acsz–27a	Aleutian–Alaska–Cascadia	199.4340	54.5960	256	15	17.94
acsz–27b	Aleutian–Alaska–Cascadia	199.6200	54.1600	256	15	5
acsz–27x	Aleutian–Alaska–Cascadia	198.9736	55.8631	256.5	15	56.24
acsz–27y	Aleutian–Alaska–Cascadia	199.1454	55.4401	256.6	15	43.82
acsz–27z	Aleutian–Alaska–Cascadia	199.3135	55.0170	256.8	15	30.88
acsz–28a	Aleutian–Alaska–Cascadia	200.8820	54.8300	253	15	17.94
acsz–28b	Aleutian–Alaska–Cascadia	201.1080	54.4000	253	15	5
acsz–28x	Aleutian–Alaska–Cascadia	200.1929	56.0559	252.5	15	56.24
acsz–28y	Aleutian–Alaska–Cascadia	200.4167	55.6406	252.7	15	43.82
acsz–28z	Aleutian–Alaska–Cascadia	200.6360	55.2249	252.9	15	30.88
acsz–29a	Aleutian–Alaska–Cascadia	202.2610	55.1330	247	15	17.94
acsz–29b	Aleutian–Alaska–Cascadia	202.5650	54.7200	247	15	5
acsz–29x	Aleutian–Alaska–Cascadia	201.2606	56.2861	245.7	15	56.24
acsz–29y	Aleutian–Alaska–Cascadia	201.5733	55.8888	246	15	43.82
acsz–29z	Aleutian–Alaska–Cascadia	201.8797	55.4908	246.2	15	30.88
acsz–30a	Aleutian–Alaska–Cascadia	203.6040	55.5090	240	15	17.94
acsz–30b acsz–30w	Aleutian–Alaska–Cascadia Aleutian–Alaska–Cascadia	203.9970	55.1200 56.9855	240	15	5
	Aleutian–Alaska–Cascadia	201.9901		239.5	15	69.12 56.24
acsz–30x acsz–30y	Aleutian–Alaska–Cascadia Aleutian–Alaska–Cascadia	202.3851 202.7724	56.6094 56.2320	239.8 240.2	15 15	56.24 43.82
acsz–30z	Aleutian–Alaska–Cascadia	203.1521	55.8534	240.2	15	30.88
acsz–30z acsz–31a	Aleutian–Alaska–Cascadia	204.8950	55.9700	236	15	17.94
acsz–31a acsz–31b	Aleutian–Alaska–Cascadia	205.3400	55.5980	236	15	5
acsz–31w	Aleutian–Alaska–Cascadia	203.0825	57.3740	234.5	15	69.12
acsz–31x	Aleutian–Alaska–Cascadia	203.5408	57.0182	234.9	15	56.24
acsz–31y	Aleutian-Alaska-Cascadia	203.9904	56.6607	235.3	15	43.82
acsz–31z	Aleutian-Alaska-Cascadia	204.4315	56.3016	235.7	15	30.88
acsz–32a	Aleutian–Alaska–Cascadia	206.2080	56.4730	236	15	17.94
acsz–32b	Aleutian–Alaska–Cascadia	206.6580	56.1000	236	15	5
acsz-32w	Aleutian-Alaska-Cascadia	204.4129	57.8908	234.3	15	69.12
acsz-32x	Aleutian-Alaska-Cascadia	204.8802	57.5358	234.7	15	56.24
acsz–32y	Aleutian-Alaska-Cascadia	205.3385	57.1792	235.1	15	43.82
acsz–32z	Aleutian-Alaska-Cascadia	205.7880	56.8210	235.5	15	30.88
acsz-33a	Aleutian-Alaska-Cascadia	207.5370	56.9750	236	15	17.94
acsz-33b	Aleutian–Alaska–Cascadia	207.9930	56.6030	236	15	5
acsz-33w	Aleutian-Alaska-Cascadia	205.7126	58.3917	234.2	15	69.12
acsz-33x	Aleutian-Alaska-Cascadia	206.1873	58.0371	234.6	15	56.24
acsz-33y	Aleutian-Alaska-Cascadia	206.6527	57.6808	235	15	43.82
acsz–33z	Aleutian–Alaska–Cascadia	207.1091	57.3227	235.4	15	30.88
acsz–34a	Aleutian–Alaska–Cascadia	208.9371	57.5124	236	15	17.94
acsz–34b	Aleutian–Alaska–Cascadia	209.4000	57.1400	236	15	5
acsz-34w	Aleutian–Alaska–Cascadia	206.9772	58.8804	233.5	15	69.12
acsz–34x	Aleutian–Alaska–Cascadia	207.4677	58.5291	233.9	15	56.24
acsz–34y	Aleutian–Alaska–Cascadia	207.9485	58.1760	234.3	15	43.82
acsz–34z	Aleutian-Alaska-Cascadia	208.4198	57.8213	234.7	15	30.88
acsz–35a	Aleutian–Alaska–Cascadia	210.2597	58.0441	230	15	17.94
acsz–35b	Aleutian–Alaska–Cascadia	210.8000	57.7000	230	15	5
acsz–35w	Aleutian–Alaska–Cascadia	208.0204	59.3199	228.8	15	69.12
acsz–35x	Aleutian–Alaska–Cascadia	208.5715	58.9906	229.3	15	56.24
acsz–35y	Aleutian–Alaska–Cascadia	209.1122	58.6590	229.7	15	43.82
acsz–35z	Aleutian–Alaska–Cascadia	209.6425	58.3252	230.2	15	30.88
acsz–36a	Aleutian–Alaska–Cascadia	211.3249	58.6565	218	15	17.94
acsz–36b	Aleutian–Alaska–Cascadia	212.0000	58.3800	218	15	5
acsz–36w	Aleutian Alaska Cascadia	208.5003	59.5894	215.6	15	69.12
acsz–36x	Aleutian Alaska-Cascadia	209.1909	59.3342	216.2	15	56.24
acsz–36y	Aleutian–Alaska–Cascadia	209.8711	59.0753	216.8	15	43.82
	Aleutian–Alaska–Cascadia	210.5412	58.8129	217.3	15	30.88
acsz–36z		212.2505	59.2720	213.7	15	17.94
acsz–37a	Aleutian–Alaska–Cascadia					-
acsz–37a acsz–37b	Aleutian–Alaska–Cascadia	212.9519	59.0312	213.7	15	5
acsz–37a						5 56.24 43.82

Table B.1 – continued

		Table B.1 - col				
Segment	Description	Longitude(°E)	Latitude(^o N)	Strike(0)	$\mathrm{Dip}(^{\mathrm{o}})$	Depth (km)
acsz–37z	Aleutian–Alaska–Cascadia	211.6079	59.5820	214.3	15	30.88
acsz–38a	Aleutian-Alaska-Cascadia	214.6555	60.1351	260.1	0	15
acsz–38b	Aleutian–Alaska–Cascadia	214.8088	59.6927	260.1	0	15
acsz–38y	Aleutian–Alaska–Cascadia	214.3737	60.9838	259	0	15
acsz–38z	Aleutian–Alaska–Cascadia	214.5362	60.5429	259	0	15
acsz–39a	Aleutian–Alaska–Cascadia	216.5607	60.2480	267	0	15
acsz–39b	Aleutian–Alaska–Cascadia	216.6068	59.7994	267	0	15
acsz–40a	Aleutian–Alaska–Cascadia	219.3069	59.7574	310.9	0	15
acsz–40b	Aleutian–Alaska–Cascadia	218.7288	59.4180	310.9	0	15
acsz–41a	Aleutian–Alaska–Cascadia	220.4832	59.3390	300.7	0	15
acsz–41b	Aleutian–Alaska–Cascadia	220.0382	58.9529	300.7	0	15
acsz–42a	Aleutian–Alaska–Cascadia	221.8835	58.9310	298.9	0	15
acsz–42b	Aleutian–Alaska–Cascadia	221.4671	58.5379	298.9	0	15
acsz–43a	Aleutian–Alaska–Cascadia	222.9711	58.6934	282.3	0	15
acsz–43b	Aleutian–Alaska–Cascadia	222.7887	58.2546	282.3	0	15
acsz–44a	Aleutian–Alaska–Cascadia	224.9379	57.9054	340.9	12	11.09
acsz–44b	Aleutian–Alaska–Cascadia	224.1596	57.7617	340.9	7	5
acsz–45a	Aleutian–Alaska–Cascadia	225.4994	57.1634	334.1	12	11.09
acsz–45b	Aleutian Alaska Cascadia	224.7740	56.9718	334.1	7	5
acsz–46a acsz–46b	Aleutian–Alaska–Cascadia Aleutian–Alaska–Cascadia	226.1459	56.3552	334.1	12 7	11.09 5
acsz–460 acsz–47a	Aleutian–Alaska–Cascadia	225.4358	56.1636	334.1 332.3	12	11.09
acsz–47a acsz–47b	Aleutian–Alaska–Cascadia	226.7731 226.0887	55.5830 55.3785	332.3	7	5
acsz–47b acsz–48a	Aleutian–Alaska–Cascadia	227.4799	54.6763	339.4	12	11.09
acsz–48b	Aleutian–Alaska–Cascadia	226.7713	54.5217	339.4	7	5
acsz–49a	Aleutian–Alaska–Cascadia	227.9482	53.8155	341.2	12	11.09
acsz–49b	Aleutian–Alaska–Cascadia	227.2462	53.6737	341.2	7	5
acsz–50a	Aleutian–Alaska–Cascadia	228.3970	53.2509	324.5	12	11.09
acsz–50b	Aleutian–Alaska–Cascadia	227.8027	52.9958	324.5	7	5
acsz–51a	Aleutian–Alaska–Cascadia	229.1844	52.6297	318.4	12	11.09
acsz–51b	Aleutian–Alaska–Cascadia	228.6470	52.3378	318.4	7	5
acsz–52a	Aleutian–Alaska–Cascadia	230.0306	52.0768	310.9	12	11.09
acsz–52b	Aleutian-Alaska-Cascadia	229.5665	51.7445	310.9	7	5
acsz–53a	Aleutian-Alaska-Cascadia	231.1735	51.5258	310.9	12	11.09
acsz–53b	Aleutian-Alaska-Cascadia	230.7150	51.1935	310.9	7	5
acsz–54a	Aleutian-Alaska-Cascadia	232.2453	50.8809	314.1	12	11.09
acsz–54b	Aleutian-Alaska-Cascadia	231.7639	50.5655	314.1	7	5
acsz–55a	Aleutian-Alaska-Cascadia	233.3066	49.9032	333.7	12	11.09
acsz–55b	Aleutian–Alaska–Cascadia	232.6975	49.7086	333.7	7	5
acsz–56a	Aleutian–Alaska–Cascadia	234.0588	49.1702	315	11	12.82
acsz–56b	Aleutian–Alaska–Cascadia	233.5849	48.8584	315	9	5
acsz–57a	Aleutian–Alaska–Cascadia	234.9041	48.2596	341	11	12.82
acsz–57b	Aleutian–Alaska–Cascadia	234.2797	48.1161	341	9	5
acsz–58a	Aleutian–Alaska–Cascadia	235.3021	47.3812	344	11	12.82
acsz–58b	Aleutian–Alaska–Cascadia	234.6776	47.2597	344	9	5
acsz–59a	Aleutian–Alaska–Cascadia	235.6432	46.5082	345	11	12.82
acsz–59b	Aleutian–Alaska–Cascadia	235.0257	46.3941	345	9	5
acsz–60a	Aleutian–Alaska–Cascadia	235.8640	45.5429	356	11	12.82
acsz–60b	Aleutian–Alaska–Cascadia	235.2363	45.5121	356	9	5
acsz–61a	Aleutian–Alaska–Cascadia	235.9106	44.6227	359	11	12.82
acsz–61b	Aleutian–Alaska–Cascadia	235.2913	44.6150	359	9	5
acsz–62a	Aleutian Alaska Cascadia	235.9229	43.7245	359	11	12.82
acsz–62b	Aleutian–Alaska–Cascadia	235.3130	43.7168	359	9	5
acsz-63a	Aleutian–Alaska–Cascadia Aleutian–Alaska–Cascadia	236.0220	42.9020	350 350	11 9	12.82
acsz–63b		235.4300	42.8254	350 345		5 12.82
acsz–64a	Aleutian–Alaska–Cascadia	235.9638	41.9818	345	11	12.82
acsz–64b	Aleutian–Alaska–Cascadia Aleutian–Alaska–Cascadia	235.3919 236.2643	41.8677	345	9	5
acsz–65a	Aleutian–Alaska–Cascadia Aleutian–Alaska–Cascadia		41.1141	345 345	11 9	12.82
acsz–65b acsz–238a	Aleutian–Alaska–Cascadia Aleutian–Alaska–Cascadia	235.7000 213.2878	41.0000	345 236.8	9 15	5 17.94
acsz–238y	Aleutian–Alaska–Cascadia	213.2676	59.8406 60.5664	236.8	15 15	43.82
acsz–238z	Aleutian–Alaska–Cascadia	212.3424 212.8119	60.2035	236.8	15 15	30.88
acsz=230Z	mounan-maska-Cascauld	212.0113	00.2033	250.0	13	30.00








Figure B.2: Central and South America Subduction Zone unit sources.

 $\label{thm:control} \textbf{Table B.2: Earthquake parameters for Central and South America Subduction Zone unit sources.}$

Segment	Description	Longitude(°E)	Latitude(^o N)	Strike(0)	Dip(°)	Depth (km)
cssz-1a	Central and South America	254.4573	20.8170	359	19	15.4
cssz–1b	Central and South America	254.0035	20.8094	359	12	5
cssz-1z	Central and South America	254.7664	20.8222	359	50	31.67
cssz-2a	Central and South America	254.5765	20.2806	336.8	19	15.4
cssz–2b	Central and South America	254.1607	20.1130	336.8	12	5
cssz–3a	Central and South America	254.8789	19.8923	310.6	18.31	15.27
cssz–3b	Central and South America	254.5841	19.5685	310.6	11.85	5
cssz–4a	Central and South America	255.6167	19.2649	313.4	17.62	15.12
cssz–4b	Central and South America	255.3056	18.9537	313.4	11.68	5
cssz–5a	Central and South America	256.2240	18.8148	302.7	16.92	15
cssz–5b	Central and South America	255.9790	18.4532	302.7	11.54	5
cssz–6a	Central and South America	256.9425	18.4383	295.1	16.23	14.87
cssz–6b cssz–7a	Central and South America Central and South America	256.7495 257.8137	18.0479 18.0339	295.1 296.9	11.38 15.54	5 14.74
cssz–7a	Central and South America	257.6079	17.6480	296.9	11.23	5
cssz–76	Central and South America	258.5779	17.7151	290.4	14.85	14.61
cssz–8b	Central and South America	258.4191	17.3082	290.4	11.08	5
cssz-9a	Central and South America	259.4578	17.4024	290.5	14.15	14.47
cssz-9b	Central and South America	259.2983	16.9944	290.5	10.92	5
cssz–10a	Central and South America	260.3385	17.0861	290.8	13.46	14.34
cssz-10b	Central and South America	260.1768	16.6776	290.8	10.77	5
cssz-11a	Central and South America	261.2255	16.7554	291.8	12.77	14.21
cssz-11b	Central and South America	261.0556	16.3487	291.8	10.62	5
cssz-12a	Central and South America	262.0561	16.4603	288.9	12.08	14.08
cssz–12b	Central and South America	261.9082	16.0447	288.9	10.46	5
cssz–13a	Central and South America	262.8638	16.2381	283.2	11.38	13.95
cssz–13b	Central and South America	262.7593	15.8094	283.2	10.31	5
cssz–14a	Central and South America	263.6066	16.1435	272.1	10.69	13.81
cssz–14b	Central and South America	263.5901	15.7024	272.1	10.15	5
cssz–15a cssz–15b	Central and South America Central and South America	264.8259 264.6462	15.8829 15.4758	293 293	10 10	13.68 5
cssz–15b	Central and South America	265.1865	16.6971	293	10	31.05
cssz–15y	Central and South America	265.0060	16.2900	293	10	22.36
cssz–16a	Central and South America	265.7928	15.3507	304.9	15	15.82
cssz–16b	Central and South America	265.5353	14.9951	304.9	12.5	5
cssz-16y	Central and South America	266.3092	16.0619	304.9	15	41.7
cssz–16z	Central and South America	266.0508	15.7063	304.9	15	28.76
cssz-17a	Central and South America	266.4947	14.9019	299.5	20	17.94
cssz–17b	Central and South America	266.2797	14.5346	299.5	15	5
cssz–17y	Central and South America	266.9259	15.6365	299.5	20	52.14
cssz–17z	Central and South America	266.7101	15.2692	299.5	20	35.04
cssz–18a	Central and South America	267.2827	14.4768	298	21.5	17.94
cssz–18b	Central and South America	267.0802	14.1078	298	15	5
cssz–18y	Central and South America	267.6888	15.2148	298	21.5	54.59
cssz–18z	Central and South America	267.4856	14.8458	298	21.5	36.27
cssz–19a	Central and South America	268.0919	14.0560	297.6	23 15	17.94
cssz–19b cssz–19y	Central and South America Central and South America	267.8943 268.4880	13.6897 14.7886	297.6 297.6	15 23	5 57.01
cssz–19y cssz–19z	Central and South America	268.2898	14.4223	297.6	23	37.48
cssz–19z cssz–20a	Central and South America	268.8929	13.6558	296.2	23 24	17.94
cssz–20a cssz–20b	Central and South America	268.7064	13.2877	296.2	15	5
cssz-20y	Central and South America	269.1796	14.2206	296.2	45.5	73.94
cssz–20z	Central and South America	269.0362	13.9382	296.2	45.5	38.28
cssz–21a	Central and South America	269.6797	13.3031	292.6	25	17.94
cssz-21b	Central and South America	269.5187	12.9274	292.6	15	5
cssz-21x	Central and South America	269.8797	13.7690	292.6	68	131.8
cssz-21y	Central and South America	269.8130	13.6137	292.6	68	85.43
cssz–21z	Central and South America	269.7463	13.4584	292.6	68	39.07
cssz–22a	Central and South America	270.4823	13.0079	288.6	25	17.94
cssz–22b	Central and South America	270.3492	12.6221	288.6	15	5
					Continued	on next page

Table B.2 – continued

		Table B.2 – con	tinued			
Segment	Description	Longitude(^o E)	Latitude(^o N)	Strike(0)	Dip(°)	Depth (km)
cssz-22x	Central and South America	270.6476	13.4864	288.6	68	131.8
cssz–22y	Central and South America	270.5925	13.3269	288.6	68	85.43
cssz–22z	Central and South America	270.5374	13.1674	288.6	68	39.07
cssz-23a	Central and South America	271.3961	12.6734	292.4	25	17.94
cssz–23b cssz–23x	Central and South America Central and South America	271.2369	12.2972	292.4	15 68	5
cssz–23y	Central and South America	271.5938 271.5279	13.1399 12.9844	292.4 292.4	68	131.8 85.43
cssz–23y	Central and South America	271.4620	12.8289	292.4	68	39.07
cssz-24a	Central and South America	272.3203	12.2251	300.2	25	17.94
cssz–24b	Central and South America	272.1107	11.8734	300.2	15	5
cssz-24x	Central and South America	272.5917	12.6799	300.2	67	131.1
cssz–24y	Central and South America	272.5012	12.5283	300.2	67	85.1
cssz–24z	Central and South America	272.4107	12.3767	300.2	67	39.07
cssz–25a	Central and South America	273.2075	11.5684	313.8	25	17.94
cssz–25b	Central and South America	272.9200	11.2746	313.8	15	5
cssz–25x	Central and South America	273.5950	11.9641	313.8	66	130.4
cssz–25y cssz–25z	Central and South America Central and South America	273.4658 273.3366	11.8322 11.7003	313.8 313.8	66 66	84.75 39.07
cssz–25z cssz–26a	Central and South America	273.8943	10.8402	320.4	25	17.94
cssz-26b	Central and South America	273.5750	10.5808	320.4	15	5
cssz–26x	Central and South America	274.3246	11.1894	320.4	66	130.4
cssz–26y	Central and South America	274.1811	11.0730	320.4	66	84.75
cssz–26z	Central and South America	274.0377	10.9566	320.4	66	39.07
cssz–27a	Central and South America	274.4569	10.2177	316.1	25	17.94
cssz–27b	Central and South America	274.1590	9.9354	316.1	15	5
cssz–27z	Central and South America	274.5907	10.3444	316.1	66	39.07
cssz–28a	Central and South America	274.9586	9.8695	297.1	22	14.54
cssz–28b cssz–28z	Central and South America Central and South America	274.7661	9.4988	297.1	11	5
cssz–26z cssz–29a	Central and South America	275.1118 275.7686	10.1643 9.4789	297.1 296.6	42.5 19	33.27 11.09
cssz–29b	Central and South America	275.5759	9.0992	296.6	7	5
cssz-30a	Central and South America	276.6346	8.9973	302.2	19	9.36
cssz–30b	Central and South America	276.4053	8.6381	302.2	5	5
cssz–31a	Central and South America	277.4554	8.4152	309.1	19	7.62
cssz–31b	Central and South America	277.1851	8.0854	309.1	3	5
cssz–31z	Central and South America	277.7260	8.7450	309.1	19	23.9
cssz–32a	Central and South America	278.1112	7.9425	303	18.67	8.49
cssz–32b	Central and South America	277.8775	7.5855	303	4	5
cssz–32z	Central and South America	278.3407	8.2927	303	21.67	24.49
cssz–33a cssz–33b	Central and South America Central and South America	278.7082 278.5785	7.6620 7.2555	287.6 287.6	18.33 6	10.23 5
cssz–33z	Central and South America	278.8328	8.0522	287.6	24.33	25.95
cssz–34a	Central and South America	279.3184	7.5592	269.5	18	17.94
cssz–34b	Central and South America	279.3223	7.1320	269.5	15	5
cssz–35a	Central and South America	280.0039	7.6543	255.9	17.67	14.54
cssz–35b	Central and South America	280.1090	7.2392	255.9	11	5
cssz–35x	Central and South America	279.7156	8.7898	255.9	29.67	79.22
cssz–35y	Central and South America	279.8118	8.4113	255.9	29.67	54.47
cssz–35z	Central and South America	279.9079	8.0328	255.9	29.67	29.72
cssz–36a	Central and South America	281.2882	7.6778	282.5	17.33	11.09
cssz–36b	Central and South America	281.1948	7.2592	282.5	7	5 70.47
cssz–36x cssz–36y	Central and South America Central and South America	281.5368 281.4539	8.7896 8.4190	282.5 282.5	32.33 32.33	79.47 52.73
cssz–36z	Central and South America	281.3710	8.0484	282.5 282.5	32.33	25.99
cssz–37a	Central and South America	282.5252	6.8289	326.9	17	10.23
cssz-37b	Central and South America	282.1629	6.5944	326.9	6	5
cssz–38a	Central and South America	282.9469	5.5973	355.4	17	10.23
cssz-38b	Central and South America	282.5167	5.5626	355.4	6	5
cssz-39a	Central and South America	282.7236	4.3108	24.13	17	10.23
cssz–39b	Central and South America	282.3305	4.4864	24.13	6	5
cssz–39z	Central and South America	283.0603	4.1604	24.13	35	24.85
cssz–40a	Central and South America	282.1940	3.3863	35.28	17	10.23
					Continued	on next page

Table B.2 - continued

Table B.2 – continued							
Segment	Description	Longitude(^o E)	Latitude(^o N)	Strike(0)	Dip(o)	Depth (kn	
cssz–40b	Central and South America	281.8427	3.6344	35.28	6	5	
cssz–40y	Central and South America	282.7956	2.9613	35.28	35	53.52	
cssz–40z	Central and South America	282.4948	3.1738	35.28	35	24.85	
cssz–41a	Central and South America	281.6890	2.6611	34.27	17	10.23	
cssz-41b	Central and South America	281.3336	2.9030	34.27	6	5	
cssz–41z	Central and South America	281.9933	2.4539	34.27	35	24.85	
cssz–42a	Central and South America	281.2266	1.9444	31.29	17	10.23	
cssz–42b	Central and South America	280.8593	2.1675	31.29	6	5	
cssz–42z	Central and South America	281.5411	1.7533	31.29	35	24.85	
cssz–43a	Central and South America	280.7297	1.1593	33.3	17	10.23	
cssz–43b	Central and South America	280.3706	1.3951	33.3	6	5	
cssz–43z	Central and South America	281.0373	0.9573	33.3	35	24.85	
cssz–44a	Central and South America	280.3018	0.4491	28.8	17	10.23	
cssz–44b	Central and South America	279.9254	0.6560	28.8	6	5	
cssz–45a	Central and South America			26.91	10	8.49	
		279.9083	-0.3259				
cssz–45b	Central and South America	279.5139	-0.1257	26.91	4	5	
cssz–46a	Central and South America	279.6461	-0.9975	15.76	10	8.49	
cssz–46b	Central and South America	279.2203	-0.8774	15.76	4	5	
cssz–47a	Central and South America	279.4972	-1.7407	6.9	10	8.49	
cssz–47b	Central and South America	279.0579	-1.6876	6.9	4	5	
cssz–48a	Central and South America	279.3695	-2.6622	8.96	10	8.49	
cssz–48b	Central and South America	278.9321	-2.5933	8.96	4	5	
cssz–48y	Central and South America	280.2444	-2.8000	8.96	10	25.85	
cssz–48z	Central and South America	279.8070	-2.7311	8.96	10	17.17	
cssz–49a	Central and South America	279.1852	-3.6070	13.15	10	8.49	
cssz–49b	Central and South America	278.7536	-3.5064	13.15	4	5	
cssz–49y	Central and South America	280.0486	-3.8082	13.15	10	25.85	
cssz-49z	Central and South America	279.6169	-3.7076	13.15	10	17.17	
cssz-50a	Central and South America	279.0652	-4.3635	4.78	10.33	9.64	
cssz–50b	Central and South America	278.6235	-4.3267	4.78	5.33	5	
cssz–51a	Central and South America	279.0349	-5.1773	359.4	10.67	10.81	
cssz–51b	Central and South America	278.5915	-5.1817	359.4	6.67	5	
cssz-52a	Central and South America	279.1047	-5.9196	349.8	11	11.96	
cssz–52b	Central and South America	278.6685	-5.9981	349.8	8	5	
cssz-53a	Central and South America	279.3044	-6.6242	339.2	10.25	11.74	
cssz–53b	Central and South America	278.8884	-6.7811	339.2	7.75	5	
cssz–53y	Central and South America	280.1024	-6.3232	339.2	19.25	37.12	
cssz–53z	Central and South America	279.7035	-6.4737	339.2	19.25	20.64	
cssz–54a	Central and South America	279.6256	-7.4907	340.8	9.5	11.53	
cssz–54b	Central and South America	279.2036	-7.6365	340.8	7.5	5	
cssz-54y	Central and South America	280.4267	-7.2137	340.8	20.5	37.29	
cssz–54z	Central and South America	280.0262	-7.3522	340.8	20.5	19.78	
	Central and South America						
cssz–55a		279.9348	-8.2452 9.4301	335.4	8.75 7.75	11.74	
cssz–55b	Central and South America	279.5269	-8.4301	335.4	7.75	5 56.4	
cssz–55x	Central and South America	281.0837	-7.7238 7.9976	335.4	21.75	56.4	
cssz–55y	Central and South America	280.7009	-7.8976	335.4	21.75	37.88	
cssz–55z	Central and South America	280.3180	-8.0714	335.4	21.75	19.35	
cssz–56a	Central and South America	280.3172	-8.9958	331.6	8	11.09	
cssz–56b	Central and South America	279.9209	-9.2072	331.6	7	5	
cssz–56x	Central and South America	281.4212	-8.4063	331.6	23	57.13	
cssz–56y	Central and South America	281.0534	-8.6028	331.6	23	37.59	
cssz–56z	Central and South America	280.6854	-8.7993	331.6	23	18.05	
cssz–57a	Central and South America	280.7492	-9.7356	328.7	8.6	10.75	
cssz–57b	Central and South America	280.3640	-9.9663	328.7	6.6	5	
cssz-57x	Central and South America	281.8205	-9.0933	328.7	23.4	57.94	
cssz-57y	Central and South America	281.4636	-9.3074	328.7	23.4	38.08	
cssz–57z	Central and South America	281.1065	-9.5215	328.7	23.4	18.22	
cssz–58a	Central and South America	281.2275	-10.5350	330.5	9.2	10.4	
cssz–58b	Central and South America	280.8348	-10.7532	330.5	6.2	5	
		281.9548	-10.1306	330.5	23.8	38.57	
cssz-58v	Central and South America	201.3340					
cssz–58y cssz–58z	Central and South America Central and South America						
cssz–58y cssz–58z cssz–59a	Central and South America Central and South America Central and South America	281.5913 281.6735	-10.3328 -11.2430	330.5 326.2	23.8 9.8	18.39 10.05	

Table B.2 - continued

Table B.2 – continued							
Segment	Description	Longitude(°E)	Latitude(^o N)	Strike(o)	Dip(o)	Depth (km)	
cssz–59b	Central and South America	281.2982	-11.4890	326.2	5.8	5	
cssz–59y	Central and South America	282.3675	-10.7876	326.2	24.2	39.06	
cssz–59z	Central and South America	282.0206	-11.0153	326.2	24.2	18.56	
cssz-60a	Central and South America	282.1864	-11.9946	326.5	10.4	9.71	
cssz-60b	Central and South America	281.8096	-12.2384	326.5	5.4	5	
cssz-60y	Central and South America	282.8821	-11.5438	326.5	24.6	39.55	
cssz–60z	Central and South America	282.5344	-11.7692	326.5	24.6	18.73	
cssz–61a	Central and South America	282.6944	-12.7263	325.5	11	9.36	
cssz–61b	Central and South America	282.3218	-12.9762	325.5	5	5	
cssz-61y	Central and South America	283.3814	-12.2649	325.5	25	40.03	
cssz-61z	Central and South America	283.0381	-12.4956	325.5	25	18.9	
cssz–62a	Central and South America	283.1980	-13.3556	319	11	9.79	
cssz–62b	Central and South America	282.8560	-13.6451	319	5.5	5	
cssz–62y	Central and South America	283.8178	-12.8300	319	27	42.03	
cssz–62z	Central and South America	283.5081	-13.0928	319	27	19.33	
cssz-63a	Central and South America	283.8032	-14.0147	317.9	11	10.23	
cssz–63b	Central and South America	283.4661	-14.3106	317.9	6	5	
cssz–63z	Central and South America	284.1032	-13.7511	317.9	29	19.77	
cssz–64a	Central and South America	284.4144	-14.6482	315.7	13	11.96	
cssz–64b	Central and South America	284.0905	-14.9540	315.7	8	5	
cssz–65a	Central and South America	285.0493	-15.2554	313.2	15	13.68	
cssz–65b	Central and South America	284.7411	-15.5715	313.2	10	5	
cssz–66a	Central and South America	285.6954	-15.7816	307.7	14.5	13.68	
cssz–66b	Central and South America	285.4190	-16.1258	307.7	10	5	
cssz–67a	Central and South America	286.4127	-16.2781	304.3	14	13.68	
cssz–67b	Central and South America	286.1566	-16.6381	304.3	10	5	
cssz–67z	Central and South America	286.6552	-15.9365	304.3	23	25.78	
cssz-68a	Central and South America	287.2481	-16.9016	311.8	14	13.68	
cssz-68b	Central and South America	286.9442	-17.2264	311.8	10	5	
cssz-68z	Central and South America	287.5291	-16.6007	311.8	26	25.78	
cssz-69a	Central and South America	287.9724	-17.5502	314.9	14	13.68	
cssz–69b	Central and South America	287.6496	-17.8590	314.9	10	5	
cssz-69y	Central and South America	288.5530	-16.9934	314.9	29	50.02	
cssz–69z	Central and South America	288.2629	-17.2718	314.9	29	25.78	
cssz–70a	Central and South America	288.6731	-18.2747	320.4	14	13.25	
cssz–70b	Central and South America	288.3193	-18.5527	320.4	9.5	5	
cssz-70y	Central and South America	289.3032	-17.7785	320.4	30	50.35	
cssz-70z	Central and South America	288.9884	-18.0266	320.4	30	25.35	
cssz-71a	Central and South America	289.3089	-19.1854	333.2	14	12.82	
cssz-71b	Central and South America	288.8968	-19.3820	333.2	9	5	
cssz–71y	Central and South America	290.0357	-18.8382	333.2	31	50.67	
cssz–71y	Central and South America				31	24.92	
	Central and South America	289.6725	-19.0118	333.2			
cssz–72a		289.6857	-20.3117	352.4	14	12.54	
cssz–72b	Central and South America	289.2250	-20.3694	352.4	8.67	5	
cssz–72z	Central and South America	290.0882	-20.2613	352.4	32	24.63	
cssz–73a	Central and South America	289.7731	-21.3061	358.9	14	12.24	
cssz–73b	Central and South America	289.3053	-21.3142	358.9	8.33	5	
cssz-73z	Central and South America	290.1768	-21.2991	358.9	33	24.34	
cssz–74a	Central and South America	289.7610	-22.2671	3.06	14	11.96	
cssz–74b	Central and South America	289.2909	-22.2438	3.06	8	5	
cssz–75a	Central and South America	289.6982	-23.1903	4.83	14.09	11.96	
cssz–75b	Central and South America	289.2261	-23.1536	4.83	8	5	
cssz-76a	Central and South America	289.6237	-24.0831	4.67	14.18	11.96	
cssz-76b	Central and South America	289.1484	-24.0476	4.67	8	5	
cssz–77a	Central and South America	289.5538	-24.9729	4.3	14.27	11.96	
cssz–77b	Central and South America	289.0750	-24.9403	4.3	8	5	
cssz–78a	Central and South America	289.4904	-25.8621	3.86	14.36	11.96	
cssz–78b	Central and South America	289.0081	-25.8328	3.86	8	5	
cssz-79a	Central and South America	289.3491	-26.8644	11.34	14.45	11.96	
cssz–79b	Central and South America	288.8712	-26.7789	11.34	8	5	
	Continuation of the fill of the control of the cont	200.0712	20.1103	11.54	U	J	
	Central and South America	280 1221	-27 7826	14 16	1/15/	11 06	
cssz–80a cssz–80b	Central and South America Central and South America	289.1231 288.6469	-27.7826 -27.6762	14.16 14.16	14.54 8	11.96 5	

Table B.2 – continued

		Table B.2 – con	tinued			
Segment	Description	Longitude(^o E)	Latitude(^o N)	Strike(o)	Dip(o)	Depth (km)
cssz–81a	Central and South America	288.8943	-28.6409	13.19	14.63	11.96
cssz–81b	Central and South America	288.4124	-28.5417	13.19	8	5
cssz–82a	Central and South America	288.7113	-29.4680	9.68	14.72	11.96
cssz–82b	Central and South America	288.2196	-29.3950	9.68	8	5
cssz–83a	Central and South America	288.5944	-30.2923	5.36	14.81	11.96
cssz–83b	Central and South America	288.0938	-30.2517	5.36	8	5
cssz–84a	Central and South America	288.5223	-31.1639	3.8	14.9	11.96
cssz–84b cssz–85a	Central and South America Central and South America	288.0163	-31.1351	3.8 2.55	8 15	5 11.96
cssz–65a	Central and South America	288.4748 287.9635	-32.0416 -32.0223	2.55	8	11.90
cssz–86a	Central and South America	288.3901	-33.0041	7.01	15	11.96
cssz–86b	Central and South America	287.8768	-32.9512	7.01	8	5
cssz–87a	Central and South America	288.1050	-34.0583	19.4	15	11.96
cssz–87b	Central and South America	287.6115	-33.9142	19.4	8	5
cssz–88a	Central and South America	287.5309	-35.0437	32.81	15	11.96
cssz-88b	Central and South America	287.0862	-34.8086	32.81	8	5
cssz-88z	Central and South America	287.9308	-35.2545	32.81	30	24.9
cssz–89a	Central and South America	287.2380	-35.5993	14.52	16.67	11.96
cssz–89b	Central and South America	286.7261	-35.4914	14.52	8	5
cssz–89z	Central and South America	287.7014	-35.6968	14.52	30	26.3
cssz–90a	Central and South America	286.8442	-36.5645	22.64	18.33	11.96
cssz–90b	Central and South America	286.3548	-36.4004	22.64	8	5
cssz–90z	Central and South America	287.2916	-36.7142	22.64	30	27.68
cssz–91a cssz–91b	Central and South America Central and South America	286.5925	-37.2488	10.9	20 8	11.96 5
cssz–91b	Central and South America	286.0721 287.0726	-37.1690 -37.3224	10.9 10.9	8 30	29.06
cssz–91z cssz–92a	Central and South America	286.4254	-38.0945	8.23	20	11.96
cssz–92b	Central and South America	285.8948	-38.0341	8.23	8	5
cssz–92z	Central and South America	286.9303	-38.1520	8.23	26.67	29.06
cssz–93a	Central and South America	286.2047	-39.0535	13.46	20	11.96
cssz-93b	Central and South America	285.6765	-38.9553	13.46	8	5
cssz-93z	Central and South America	286.7216	-39.1495	13.46	23.33	29.06
cssz–94a	Central and South America	286.0772	-39.7883	3.4	20	11.96
cssz–94b	Central and South America	285.5290	-39.7633	3.4	8	5
cssz–94z	Central and South America	286.6255	-39.8133	3.4	20	29.06
cssz–95a	Central and South America	285.9426	-40.7760	9.84	20	11.96
cssz–95b	Central and South America	285.3937	-40.7039	9.84	8	5
cssz–95z cssz–96a	Central and South America Central and South America	286.4921 285.7839	-40.8481	9.84 7.6	20 20	29.06 11.96
cssz–96b	Central and South America	285.2245	-41.6303 -41.5745	7.6	8	5
cssz-96x	Central and South America	287.4652	-41.7977	7.6	20	63.26
cssz–96y	Central and South America	286.9043	-41.7419	7.6	20	46.16
cssz–96z	Central and South America	286.3439	-41.6861	7.6	20	29.06
cssz-97a	Central and South America	285.6695	-42.4882	5.3	20	11.96
cssz–97b	Central and South America	285.0998	-42.4492	5.3	8	5
cssz–97x	Central and South America	287.3809	-42.6052	5.3	20	63.26
cssz–97y	Central and South America	286.8101	-42.5662	5.3	20	46.16
cssz–97z	Central and South America	286.2396	-42.5272	5.3	20	29.06
cssz–98a	Central and South America	285.5035	-43.4553	10.53	20	11.96
cssz–98b	Central and South America	284.9322	-43.3782	10.53	8	5
cssz–98x	Central and South America	287.2218	-43.6866	10.53	20	63.26
cssz–98y cssz–98z	Central and South America Central and South America	286.6483 286.0755	-43.6095 -43.5324	10.53	20 20	46.16
cssz–96z cssz–99a	Central and South America	285.3700	-43.5324 -44.2595	10.53 4.86	20 20	29.06 11.96
cssz–99b	Central and South America	284.7830	-44.2237	4.86	8	5
cssz–99x	Central and South America	287.1332	-44.3669	4.86	20	63.26
cssz-99y	Central and South America	286.5451	-44.3311	4.86	20	46.16
cssz–99z	Central and South America	285.9574	-44.2953	4.86	20	29.06
cssz-100a	Central and South America	285.2713	-45.1664	5.68	20	11.96
cssz-100b	Central and South America	284.6758	-45.1246	5.68	8	5
cssz-100x	Central and South America	287.0603	-45.2918	5.68	20	63.26
cssz-100y	Central and South America	286.4635	-45.2500	5.68	20	46.16
					Continued	on next page

Table B.2 – continued

		Table B.2 – con	itinued			
Segment	Description	Longitude(^o E)	Latitude(^o N)	Strike(0)	Dip(o)	Depth (km)
cssz-100z	Central and South America	285.8672	-45.2082	5.68	20	29.06
cssz-101a	Central and South America	285.3080	-45.8607	352.6	20	9.36
cssz-101b	Central and South America	284.7067	-45.9152	352.6	5	5
cssz-101y	Central and South America	286.5089	-45.7517	352.6	20	43.56
cssz-101z	Central and South America	285.9088	-45.8062	352.6	20	26.46
cssz-102a	Central and South America	285.2028	-47.1185	17.72	5	9.36
cssz-102b	Central and South America	284.5772	-46.9823	17.72	5	5
cssz-102y	Central and South America	286.4588	-47.3909	17.72	5	18.07
cssz-102z	Central and South America	285.8300	-47.2547	17.72	5	13.72
cssz–103a	Central and South America	284.7075	-48.0396	23.37	7.5	11.53
cssz–103b	Central and South America	284.0972	-47.8630	23.37	7.5	5
cssz–103x	Central and South America	286.5511	-48.5694	23.37	7.5	31.11
cssz-103y	Central and South America	285.9344	-48.3928	23.37	7.5	24.58
cssz–103z	Central and South America	285.3199	-48.2162	23.37	7.5	18.05
cssz-104a	Central and South America	284.3440	-48.7597	14.87	10	13.68
cssz–104b	Central and South America	283.6962	-48.6462	14.87	10	5
cssz–104x	Central and South America	286.2962	-49.1002	14.87	10	39.73
cssz-104y	Central and South America	285.6440	-48.9867	14.87	10	31.05
cssz-104z	Central and South America Central and South America	284.9933	-48.8732	14.87	10	22.36
cssz-105a	Central and South America	284.2312 283.5518	-49.4198	0.25	9.67 9.67	13.4
cssz–105b cssz–105x	Central and South America	286.2718	-49.4179 -49.4255	0.25 0.25	9.67	5 38.59
cssz–105x	Central and South America	285.5908	-49.4236	0.25	9.67	30.2
cssz–105z	Central and South America	284.9114	-49.4217	0.25	9.67	21.8
cssz–106a	Central and South America	284.3730	-50.1117	347.5	9.25	13.04
cssz-106b	Central and South America	283.6974	-50.2077	347.5	9.25	5
cssz-106x	Central and South America	286.3916	-49.8238	347.5	9.25	37.15
cssz–106y	Central and South America	285.7201	-49.9198	347.5	9.25	29.11
cssz–106z	Central and South America	285.0472	-50.0157	347.5	9.25	21.07
cssz-107a	Central and South America	284.7130	-50.9714	346.5	9	12.82
cssz–107b cssz–107x	Central and South America Central and South America	284.0273 286.7611	-51.0751 -50.6603	346.5 346.5	9 9	5 36.29
cssz–107x cssz–107y	Central and South America	286.0799	-50.7640	346.5	9	28.47
cssz–107z	Central and South America	285.3972	-50.8677	346.5	9	20.64
cssz–108a	Central and South America	285.0378	-51.9370	352	8.67	12.54
cssz-108b	Central and South America	284.3241	-51.9987	352	8.67	5
cssz-108x	Central and South America	287.1729	-51.7519	352	8.67	35.15
cssz–108y	Central and South America	286.4622	-51.8136	352	8.67	27.61
cssz–108z	Central and South America	285.7505	-51.8753	352	8.67	20.07
cssz–109a cssz–109b	Central and South America Central and South America	285.2635 284.5326	-52.8439 -52.8974	353.1 353.1	8.33 8.33	12.24 5
cssz-109b	Central and South America	287.4508	-52.6834	353.1	8.33	33.97
cssz–109y	Central and South America	286.7226	-52.7369	353.1	8.33	26.73
cssz–109z	Central and South America	285.9935	-52.7904	353.1	8.33	19.49
cssz-110a	Central and South America	285.5705	-53.4139	334.2	8	11.96
cssz–110b	Central and South America	284.8972	-53.6076	334.2	8	5
cssz–110x	Central and South America	287.5724	-52.8328	334.2	8	32.83
cssz-110y	Central and South America	286.9081	-53.0265	334.2	8	25.88
cssz–110z	Central and South America	286.2408	-53.2202	334.2	8	18.92
cssz–111a cssz–111b	Central and South America Central and South America	286.1627 285.6382	-53.8749 -54.1958	313.8 313.8	8 8	11.96 5
cssz–111b	Central and South America	287.7124	-52.9122	313.8	8	32.83
cssz-111y	Central and South America	287.1997	-53.2331	313.8	8	25.88
cssz–111z	Central and South America	286.6832	-53.5540	313.8	8	18.92
cssz-112a	Central and South America	287.3287	-54.5394	316.4	8	11.96
cssz–112b	Central and South America	286.7715	-54.8462	316.4	8	5
cssz-112x	Central and South America	288.9756	-53.6190	316.4	8	32.83
cssz–112y	Central and South America	288.4307	-53.9258	316.4	8	25.88
cssz–112z	Central and South America	287.8817	-54.2326	316.4	8	18.92
cssz-113a	Central and South America	288.3409	-55.0480	307.6	8	11.96
cssz–113b cssz–113x	Central and South America Central and South America	287.8647 289.7450	-55.4002 -53.9914	307.6 307.6	8 8	5 32.83
cssz–113x	Central and South America	289.2810	-54.3436	307.6	8	25.88
cssz–113y	Central and South America	288.8130	-54.6958	307.6	8	18.92
cssz–114a	Central and South America	289.5342	-55.5026	301.5	8	11.96
					Continued	on next page

Table B.2 – continued

Segment	Description	Longitude(^o E)	Latitude(⁰ N)	Strike(0)	Dip(o)	Depth (km)
cssz–114b	Central and South America	289.1221	-55.8819	301.5	8	5
cssz-114x	Central and South America	290.7472	-54.3647	301.5	8	32.83
cssz-114y	Central and South America	290.3467	-54.7440	301.5	8	25.88
cssz-114z	Central and South America	289.9424	-55.1233	301.5	8	18.92
cssz-115a	Central and South America	290.7682	-55.8485	292.7	8	11.96
cssz-115b	Central and South America	290.4608	-56.2588	292.7	8	5
cssz-115x	Central and South America	291.6714	-54.6176	292.7	8	32.83
cssz-115y	Central and South America	291.3734	-55.0279	292.7	8	25.88
cssz–115z	Central and South America	291.0724	-55.4382	292.7	8	18.92






Figure B.3: Eastern Philippines Subduction Zone unit sources.

 $\label{eq:Basic_Basic_Basic} \textbf{Earthquake parameters for Eastern Philippines Subduction Zone unit sources.}$

Segment	Description	Longitude(^o E)	Latitude(^o N)	Strike(⁰)	Dip(o)	Depth (km)
epsz-0a	Eastern Philippines	128.5264	1.5930	180	44	26.92
epsz–0b	Eastern Philippines	128.8496	1.5930	180	26	5
epsz–1a	Eastern Philippines	128.5521	2.3289	153.6	44.2	27.62
epsz–1b	Eastern Philippines	128.8408	2.4720	153.6	26.9	5
epsz–2a	Eastern Philippines	128.1943	3.1508	151.9	45.9	32.44
epsz–2b	Eastern Philippines	128.4706	3.2979	151.9	32.8	5.35
epsz–3a	Eastern Philippines	127.8899	4.0428	155.2	57.3	40.22
epsz–3b	Eastern Philippines	128.1108	4.1445	155.2	42.7	6.31
epsz–4a	Eastern Philippines	127.6120	4.8371	146.8	71.4	48.25
epsz–4b	Eastern Philippines	127.7324	4.9155	146.8	54.8	7.39
epsz–5a	Eastern Philippines	127.3173	5.7040	162.9	79.9	57.4
epsz–5b	Eastern Philippines	127.3930	5.7272	162.9	79.4	8.25
epsz–6a	Eastern Philippines	126.6488	6.6027	178.9	48.6	45.09
epsz–6b	Eastern Philippines	126.9478	6.6085	178.9	48.6	7.58
epsz–7a	Eastern Philippines	126.6578	7.4711	175.8	50.7	45.52
epsz–7b	Eastern Philippines	126.9439	7.4921	175.8	50.7	6.83
epsz–8a	Eastern Philippines	126.6227	8.2456	163.3	56.7	45.6
epsz–8b	Eastern Philippines	126.8614	8.3164	163.3	48.9	7.92
epsz–9a	Eastern Philippines	126.2751	9.0961	164.1	47	43.59
epsz–9b	Eastern Philippines	126.5735	9.1801	164.1	44.9	8.3
epsz–10a	Eastern Philippines	125.9798	9.9559	164.5	43.1	42.25
epsz–10b	Eastern Philippines	126.3007	10.0438	164.5	43.1	8.09
epsz–11a	Eastern Philippines	125.6079	10.6557	155	37.8	38.29
epsz–11b	Eastern Philippines	125.9353	10.8059	155	37.8	7.64
epsz–12a	Eastern Philippines	125.4697	11.7452	172.1	36	37.01
epsz–12b	Eastern Philippines	125.8374	11.7949	172.1	36	7.62
epsz–13a	Eastern Philippines	125.2238	12.1670	141.5	32.4	33.87
epsz–13b	Eastern Philippines	125.5278	12.4029	141.5	32.4	7.08
epsz–14a	Eastern Philippines	124.6476	13.1365	158.2	23	25.92
epsz–14b	Eastern Philippines	125.0421	13.2898	158.2	23	6.38
epsz–15a	Eastern Philippines	124.3107	13.9453	156.1	24.1	26.51
epsz–15b	Eastern Philippines	124.6973	14.1113	156.1	24.1	6.09
epsz–16a	Eastern Philippines	123.8998	14.4025	140.3	19.5	21.69
epsz–16b	Eastern Philippines	124.2366	14.6728	140.3	19.5	5
epsz–17a	Eastern Philippines	123.4604	14.7222	117.6	15.3	18.19
epsz–17b	Eastern Philippines	123.6682	15.1062	117.6	15.3	5
epsz–18a	Eastern Philippines	123.3946	14.7462	67.4	15	17.94
epsz–18b	Eastern Philippines	123.2219	15.1467	67.4	15	5
epsz–19a	Eastern Philippines	121.3638	15.7400	189.6	15	17.94
epsz–19b	Eastern Philippines	121.8082	15.6674	189.6	15	5
epsz–20a	Eastern Philippines	121.6833	16.7930	203.3	15	17.94
epsz–20b	Eastern Philippines	122.0994	16.6216	203.3	15	5
epsz–21a	Eastern Philippines	121.8279	17.3742	184.2	15	17.94
epsz–21b	Eastern Philippines	122.2814	17.3425	184.2	15	5




Figure B.4: Kamchatka-Bering Subduction Zone unit sources.

 $\label{thm:control} \mbox{Table B.4: Earthquake parameters for Kamchatka-Bering Subduction Zone unit sources.}$

Segment	Description	Longitude(^o E)	Latitude(^o N)	Strike(0)	Dip(o)	Depth (km)
kbsz-1a	Kamchatka-Bering	g 161.8374	57.5485	201.5	29	26.13
kbsz-1b	Kamchatka-Bering	g 162.5162	57.4030	202.1	25	5
kbsz-2a	Kamchatka-Bering	g 162.4410	58.3816	201.7	29	26.13
kbsz-2b	Kamchatka-Bering	g 163.1344	58.2343	202.3	25	5
kbsz-2z	Kamchatka-Bering	g 161.7418	58.5249	201.1	29	50.37
kbsz-3a	Kamchatka-Bering	g 163.5174	59.3493	218.9	29	26.13
kbsz-3b	Kamchatka-Bering	g 164.1109	59.1001	219.4	25	5
kbsz-3z	Kamchatka-Bering	g 162.9150	59.5958	218.4	29	50.37
kbsz–4a	Kamchatka-Bering	g 164.7070	60.0632	222.2	29	26.13
kbsz–4b	Kamchatka-Bering	g 165.2833	59.7968	222.7	25	5
kbsz–4z	Kamchatka-Bering	g 164.1212	60.3270	221.7	29	50.37
kbsz-5a	Kamchatka-Bering	g 165.8652	60.7261	220.5	29	26.13
kbsz–5b	Kamchatka-Bering	g 166.4692	60.4683	221	25	5






Figure B.5: Kamchatka-Kuril-Japan-Izu-Mariana-Yap Subduction Zone unit sources.

 $\label{thm:continuous} \parbox{Table B.5: Earthquake parameters for Kamchatka-Kuril-Japan-Izu-Mariana-Yap Subduction Zone unit sources.}$

Segment	Description	Longitude(^o E)) Latitude(^o N)	Strike(0)	Dip(o)	Depth (kı
kisz–0a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap		56.3667	194.4	29	26.13
kisz–0b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	163.5057	56.2677	195	25	5
kisz–0z	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	162.1309	56.4618	193.8	29	50.37
kisz–1a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap		55.5017	195	29	26.13
kisz–1b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap		55.4000	195	25	5
kisz–1y	Kamchatka-Kuril-Japan-Izu-Mariana-Yap		55.7050	195	29	74.61
kisz–1z	Kamchatka-Kuril-Japan-Izu-Mariana-Yap		55.6033	195	29	50.37
kisz–2a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap		54.6784	200	29	26.13
kisz–2b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap		54.5440	200	25	5
kisz–2y	Kamchatka-Kuril-Japan-Izu-Mariana-Yap		54.9471	200	29	74.61
kisz–2z	Kamchatka-Kuril-Japan-Izu-Mariana-Yap		54.8127	200	29	50.37
kisz–3a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap		53.8714	204	29	26.13
kisz–3b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap		53.7116	204	25	5
kisz–3y	Kamchatka-Kuril-Japan-Izu-Mariana-Yap		54.1910	204	29	74.61
kisz–3z	Kamchatka-Kuril-Japan-Izu-Mariana-Yap		54.0312	204	29	50.37
kisz–4a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap		53.1087	210	29	26.13
kisz–4b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap		52.9123	210	25	5
kisz–4y	Kamchatka-Kuril-Japan-Izu-Mariana-Yap		53.5015	210	29	74.61
kisz–4z	Kamchatka-Kuril-Japan-Izu-Mariana-Yap		53.3051	210	29	50.37
kisz–5a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap		52.4113	218	29	26.13
kisz–5b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap		52.1694	218	25	5
kisz–5y	Kamchatka-Kuril-Japan-Izu-Mariana-Yap		52.8950	218	29	74.61
kisz–5z	Kamchatka-Kuril-Japan-Izu-Mariana-Yap		52.6531	218	29	50.37
kisz–6a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap		51.7034	218	29	26.13
kisz–6b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap		51.4615	218	25	5
kisz–6y	Kamchatka-Kuril-Japan-Izu-Mariana-Yap		52.1871	218	29	74.61
kisz–6z	Kamchatka-Kuril-Japan-Izu-Mariana-Yap		51.9452	218	29	50.37
kisz–7a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap		50.9549	214	29	26.13
kisz–7b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap		50.7352	214	25	5
kisz–7y	Kamchatka-Kuril-Japan-Izu-Mariana-Yap		51.3942	214	29	74.61
kisz–7z	Kamchatka-Kuril-Japan-Izu-Mariana-Yap		51.1745	214	29	50.37
kisz–8a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap		50.2459	218	31	27.7
kisz–8b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap		50.0089	218	27	5
kisz–8y	Kamchatka-Kuril-Japan-Izu-Mariana-Yap		50.7199	218	31	79.2
kisz–8z	Kamchatka-Kuril-Japan-Izu-Mariana-Yap		50.4829	218	31	53.45
kisz–9a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap		49.5583	220	31	27.7
kisz–9b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap		49.3109	220	27	5
kisz–9y	Kamchatka-Kuril-Japan-Izu-Mariana-Yap		50.0533	220	31	79.2
kisz–9z	Kamchatka-Kuril-Japan-Izu-Mariana-Yap		49.8058	220	31	53.45
kisz–10a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap		48.8804	221	31	27.7
kisz–10b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap		48.6278	221	27	5
kisz–10y	Kamchatka-Kuril-Japan-Izu-Mariana-Yap		49.3856	221	31	79.2
kisz–10z	Kamchatka-Kuril-Japan-Izu-Mariana-Yap		49.1330	221	31	53.45
kisz–11a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap		48.1821	219	31	27.7
kisz–11b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap		47.9398	219	27	5
kisz–11y	Kamchatka-Kuril-Japan-Izu-Mariana-Yap		48.6667	219	31	79.2
kisz–11z	Kamchatka-Kuril-Japan-Izu-Mariana-Yap		48.4244	219	31	53.45
kisz–11c	Kamchatka-Kuril-Japan-Izu-Mariana-Yap		47.5374	39	57.89	4.602
kisz–12a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap		47.4729	217	31	27.7
kisz–12b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap		47.2320	217	27	5
kisz–12y	Kamchatka-Kuril-Japan-Izu-Mariana-Yap		47.9363	217	31	79.2
kisz–12z	Kamchatka-Kuril-Japan-Izu-Mariana-Yap		47.7046	217	31	53.45
kisz–12c	Kamchatka-Kuril-Japan-Izu-Mariana-Yap		46.8473	37	57.89	4.602
kisz–13a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap		46.7564	218	31	27.7
kisz–13b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap		46.5194	218	27	5
kisz–13y	Kamchatka-Kuril-Japan-Izu-Mariana-Yap		47.2304	218	31	79.2
kisz–13z	Kamchatka-Kuril-Japan-Izu-Mariana-Yap		46.9934	218	31	53.45
kisz–13c	Kamchatka-Kuril-Japan-Izu-Mariana-Yap		46.1257	38	57.89	4.602
kisz–14a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	152.3657	46.1514	225	23	24.54
kisz–14b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	152.7855	45.8591	225	23	5

Table B.5 – continued

Table B.5 – continued						
Segment	Description I	Longitude(^o I	E) Latitude(^o N)	Strike(^o) Dip(°) I	Depth (km)
kisz–14y	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	151.5172	46.7362	225	23	63.62
kisz–14z	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	151.9426	46.4438	225	23	44.08
kisz–14c	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	153.4468	45.3976	45	57.89	4.602
kisz–15a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	151.4663	45.5963	233	25	23.73
kisz–15b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	151.8144	45.2712	233	22	5
kisz–15y	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	150.7619	46.2465	233	25	65.99
kisz–15z	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	151.1151	45.9214	233	25	44.86
kisz–16a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	150.4572	45.0977	237	25	23.73
kisz–16b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	150.7694	44.7563	237	22	5
kisz–16y	Kamchatka-Kuril-Japan-Izu-Mariana-Yap Kamchatka-Kuril-Japan-Izu-Mariana-Yap	149.8253	45.7804	237	25	65.99
kisz–16z kisz–17a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	150.1422 149.3989	45.4390 44.6084	237 237	25 25	44.86 23.73
kisz–17a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	149.7085	44.2670	237	22	23.73 5
kisz–170 kisz–17y	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	148.7723	45.2912	237	25	65.99
kisz–17z	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	149.0865	44.9498	237	25	44.86
kisz–18a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	148.3454	44.0982	235	25	23.73
kisz–18b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	148.6687	43.7647	235	22	5
kisz-18y	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	147.6915	44.7651	235	25	65.99
kisz–18z	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	148.0194	44.4316	235	25	44.86
kisz-19a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	147.3262	43.5619	233	25	23.73
kisz-19b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	147.6625	43.2368	233	22	5
kisz–19y	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	146.6463	44.2121	233	25	65.99
kisz–19z	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	146.9872	43.8870	233	25	44.86
kisz–20a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap		43.0633	237	25	23.73
kisz–20b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	146.6531	42.7219	237	22	5
kisz–20y	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	145.7410	43.7461	237	25	65.99
kisz–20z	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	146.0470	43.4047	237	25	44.86
kisz–21a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	145.3331	42.5948	239	25	23.73
kisz–21b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	145.6163	42.2459	239	22	5
kisz–21y kisz–21z	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	144.7603	43.2927	239 239	25	65.99 44.86
kisz–21z kisz–22a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap Kamchatka-Kuril-Japan-Izu-Mariana-Yap	145.0475 144.3041	42.9438 42.1631	242	25 25	23.73
kisz–22b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	144.5605	41.8037	242	22	5
kisz–22y	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	143.7854	42.8819	242	25	65.99
kisz–22z	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	144.0455	42.5225	242	25	44.86
kisz-23a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	143.2863	41.3335	202	21	21.28
kisz-23b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	143.8028	41.1764	202	19	5
kisz-23v	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	140.6816	42.1189	202	21	110.9
kisz-23w	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	141.2050	41.9618	202	21	92.95
kisz–23x	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	141.7273	41.8047	202	21	75.04
kisz–23y	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	142.2482	41.6476	202	21	57.12
kisz–23z	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	142.7679	41.4905	202	21	39.2
kisz–24a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	142.9795	40.3490	185	21	21.28
kisz–24b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	143.5273	40.3125	185	19	5
kisz–24x	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	141.3339	40.4587	185	21	75.04
kisz–24y kisz–24z	Kamchatka-Kuril-Japan-Izu-Mariana-Yap Kamchatka-Kuril-Japan-Izu-Mariana-Yap	141.8827	40.4221	185	21	57.12
kisz–24z kisz–25a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	142.4312 142.8839	40.3856 39.4541	185 185	21 21	39.2 21.28
kisz–25b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	143.4246	39.4176	185	19	5
kisz–25y	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	141.8012	39.5272	185	21	57.12
kisz–25z	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	142.3426	39.4907	185	21	39.2
kisz–26a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	142.7622	38.5837	188	21	21.28
kisz–26b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	143.2930	38.5254	188	19	5
kisz–26x	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	141.1667	38.7588	188	21	75.04
kisz-26y	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	141.6990	38.7004	188	21	57.12
kisz–26z	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	142.2308	38.6421	188	21	39.2
kisz-27a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	142.5320	37.7830	198	21	21.28
kisz–27b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	143.0357	37.6534	198	19	5
kisz–27x	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	141.0142	38.1717	198	21	75.04
kisz–27y	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	141.5210	38.0421	198	21	57.12
kisz–27z	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	142.0269	37.9126	198	21	39.2
kisz–28a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	142.1315	37.0265	208	21	21.28
				Con	tinued or	n next page

131

Table B.5 – continued

Riss-28b	
kisz-28y Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141,2016 37,4202 208 21 kisz-28z Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141,2016 37,4202 208 21 kisz-29a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141,6671 37,2234 208 21 kisz-29a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141,6671 37,2234 208 21 kisz-29x Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142,0416 36,0481 211 19 kisz-29y Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142,0416 36,0481 211 21 kisz-30a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141,0503 36,4800 211 21 kisz-30a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141,053 35,4332 205 21 kisz-30a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141,0583 35,4332 205 21 kisz-30a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 140,0883 35,6104 205 21 kisz-30b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 140,0883 36,6104 205 21 kisz-31b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 140,0895 34,4789 190 22 kisz-31b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 140,0895 34,4789 190 22 kisz-31x Kamchatka-Kuril-Japan-Izu-Mariana-Yap 138,7021 34,7682 190 22 kisz-31x Kamchatka-Kuril-Japan-Izu-Mariana-Yap 138,9012 34,7682 190 22 kisz-31z Kamchatka-Kuril-Japan-Izu-Mariana-Yap 140,1979 34,5512 190 22 kisz-32a kamchatka-Kuril-Japan-Izu-Mariana-Yap 140,1979 34,5512 190 22 kisz-33a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141,1958 33,0921 180 24,685	epth (km
kisz-28y Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141,2016 37,4202 208 21 kisz-28z Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141,2016 37,4202 208 21 kisz-29a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141,6671 37,2234 208 21 kisz-29a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141,6671 37,2234 208 21 kisz-29x Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142,0416 36,0481 211 19 kisz-29y Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142,0416 36,0481 211 21 kisz-30a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141,0503 36,4800 211 21 kisz-30a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141,053 35,4332 205 21 kisz-30a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141,0583 35,4332 205 21 kisz-30a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 140,0883 35,6104 205 21 kisz-30b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 140,0883 36,6104 205 21 kisz-31b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 140,0895 34,4789 190 22 kisz-31b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 140,0895 34,4789 190 22 kisz-31x Kamchatka-Kuril-Japan-Izu-Mariana-Yap 138,7021 34,7682 190 22 kisz-31x Kamchatka-Kuril-Japan-Izu-Mariana-Yap 138,9012 34,7682 190 22 kisz-31z Kamchatka-Kuril-Japan-Izu-Mariana-Yap 140,1979 34,5512 190 22 kisz-32a kamchatka-Kuril-Japan-Izu-Mariana-Yap 140,1979 34,5512 190 22 kisz-33a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141,1958 33,0921 180 24,685	5
kisz-28y Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141.2016 37.4202 208 21 21 21 21 22 23 23 23	75.04
kisz-29s	57.12
kisz-29a	39.2
kisz-29b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.0416 36.0481 211 19 19 12 19 12 19 12 19 14 10 10 10 11 12 13 14 15 16 16 17 17 18 18 18 18 18 18	21.28
kisz-29y Kamchatka-Kuril-Japan-Izu-Mariana-Yap 140.7029 36.6960 211 21 21 21 21 22 23 23	5
kisz-29a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141,1506 36,4800 211 21 kisz-30b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141,0553 35,4332 205 19 kisz-30b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141,5207 35,2560 205 19 kisz-30y Kamchatka-Kuril-Japan-Izu-Mariana-Yap 140,5203 35,7876 205 21 kisz-31b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 140,6956 34,4789 190 22 kisz-31v Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141,1927 34,4066 190 20 kisz-31v Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141,1927 34,4066 190 22 kisz-31v Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141,1927 34,6682 190 22 kisz-31v Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141,0594 34,6512 190 22 kisz-31z Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141,0594 33,0921 180 22 kisz-3ba Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141,5098	57.12
kisz-30a kamchatka-Kuril-Japan-Izu-Mariana-Yap 141.0523 35.2560 205 21 kisz-30y Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141.5207 35.2560 205 21 kisz-30y Kamchatka-Kuril-Japan-Izu-Mariana-Yap 140.1204 35.7876 205 21 kisz-31b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 140.5883 35.6104 205 21 kisz-31b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 140.5883 35.6104 205 21 kisz-31b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141.1927 34.4066 190 20 kisz-31v Kamchatka-Kuril-Japan-Izu-Mariana-Yap 138.2025 34.8405 190 22 kisz-31y Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141.0551 33.0921 180 22 kisz-31y Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141.0551 33.0921 180 21 kisz-32b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141.0551 33.0921 180 21 kisz-33b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141.0568	39.2
kisz-30b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141,5207 35,2560 205 19 kisz-30z Kamchatka-Kuril-Japan-Izu-Mariana-Yap 140,1204 35,7876 205 21 kisz-31a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 140,6986 34,4789 190 22 kisz-31b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141,1927 34,4066 190 20 kisz-31v Kamchatka-Kuril-Japan-Izu-Mariana-Yap 138,2025 34,8405 190 22 kisz-31v Kamchatka-Kuril-Japan-Izu-Mariana-Yap 138,2021 34,6682 190 22 kisz-31v Kamchatka-Kuril-Japan-Izu-Mariana-Yap 140,1979 34,6512 190 22 kisz-31v Kamchatka-Kuril-Japan-Izu-Mariana-Yap 140,1979 34,5512 190 22 kisz-32a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141,0594 33,0921 180 21.69 kisz-33a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141,0594 32,1473 173.8 27.65 kisz-34b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141,0589	21.28
kisz-30y Kamchatka-Kuril-Japan-Izu-Mariana-Yap 140,1204 35,7876 205 21 kisz-30z Kamchatka-Kuril-Japan-Izu-Mariana-Yap 140,5883 35,6104 205 21 kisz-31b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141,1927 34,4066 190 20 kisz-31b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141,1927 34,4066 190 20 kisz-31w Kamchatka-Kuril-Japan-Izu-Mariana-Yap 138,2025 34,8405 190 22 kisz-31y Kamchatka-Kuril-Japan-Izu-Mariana-Yap 139,2012 34,6958 190 22 kisz-31y Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141,0551 33,0921 180 22 kisz-32b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141,0591 33,0921 180 21,69 kisz-32b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141,5968 33,0921 180 21,69 kisz-34a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141,5968 31,1851 172,1 25 kisz-35b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141,4154	5
kisz-30z Kamchatka-Kuril-Japan-Izu-Mariana-Yap 440.5883 35.6104 205 21 kisz-31b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 140.6956 34.4789 190 22 kisz-31b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141.1927 34.4066 190 20 kisz-31v Kamchatka-Kuril-Japan-Izu-Mariana-Yap 138.2025 34.8405 190 22 kisz-31v Kamchatka-Kuril-Japan-Izu-Mariana-Yap 138.2021 34.6958 190 22 kisz-31y Kamchatka-Kuril-Japan-Izu-Mariana-Yap 140.1979 34.6512 190 22 kisz-31y Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141.0551 33.0921 180 32 kisz-32a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141.0551 33.0921 180 21.69 kisz-33a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141.5998 32.1473 173.8 18.27.65 kisz-34b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141.1869 31.1851 172.1 25 kisz-35b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141.0626<	57.12
kisz-31a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 140.6956 34.4788 190 22 kisz-31b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141.1927 34.4066 190 20 kisz-31w Kamchatka-Kuril-Japan-Izu-Mariana-Yap 138.7021 34.7682 190 22 kisz-31y Kamchatka-Kuril-Japan-Izu-Mariana-Yap 139.2012 34.6958 190 22 kisz-31z Kamchatka-Kuril-Japan-Izu-Mariana-Yap 140.1979 34.5512 190 22 kisz-31z Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141.0979 34.5512 190 22 kisz-32a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141.0979 34.5512 190 22 kisz-33a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141.0924 32.1047 173.8 27.65 kisz-33b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141.1869 31.1851 172.1 25 kisz-35a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141.6585 31.2408 172.1 15.38 kisz-35b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141.0581 </td <td>39.2</td>	39.2
kisz-31v Kamchatka-Kuril-Japan-Izu-Mariana-Yap 138.2025 34.8405 190 22 kisz-31w Kamchatka-Kuril-Japan-Izu-Mariana-Yap 138.7021 34.7682 190 22 kisz-31y Kamchatka-Kuril-Japan-Izu-Mariana-Yap 139.2012 34.6935 190 22 kisz-31z Kamchatka-Kuril-Japan-Izu-Mariana-Yap 140.1979 34.5512 190 22 kisz-32a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141.0551 33.0921 180 32 kisz-32b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141.5098 33.0921 180 21.69 kisz-33b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141.5869 31.1851 172.1 25 kisz-34b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141.6863 31.2408 172.1 25 kisz-35b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141.661 30.2899 163 14.03 kisz-36a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.0670 29.4012 161.7 25.73 kisz-37a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.0670	22.1
kisz-31w Kamchatka-Kuril-Japan-Izu-Mariana-Yap 138.7021 34.7682 190 22 kisz-31y Kamchatka-Kuril-Japan-Izu-Mariana-Yap 139.2012 34.6938 190 22 kisz-31z Kamchatka-Kuril-Japan-Izu-Mariana-Yap 140.1979 34.5512 190 22 kisz-32a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141.0551 33.0921 180 32 kisz-32a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141.0551 33.0921 180 32 kisz-33b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141.5986 33.0921 180 22 kisz-33b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141.5966 32.1473 173.8 18.27 kisz-34a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141.1869 31.1831 172.1 15.38 kisz-35a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141.9662 30.2899 163 14.03 kisz-35b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.0670 29.4012 161.7 25.73 kisz-36b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.	5
kisz-31x Kamchatka-Kuril-Japan-Izu-Mariana-Yap 139.2012 34,6958 190 22 kisz-31y Kamchatka-Kuril-Japan-Izu-Mariana-Yap 139,6997 34,6235 190 22 kisz-32a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 140,1979 34,5512 190 22 kisz-32b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141,5098 33,0921 180 32 kisz-33b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141,5098 33,0921 180 21,69 kisz-33b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141,5596 32,1473 173.8 182,7 kisz-34b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141,6585 31,2408 172,1 25 kisz-35a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141,6662 30,2899 163 140,3 kisz-35b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142,0670 29,4012 161,7 25,73 kisz-36b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142,0210 28,3322 154,7 20 kisz-37a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 14	115.8
kisz-31y Kamchatka-Kuril-Japan-Izu-Mariana-Yap 139.6997 34.6235 190 22 kisz-31z Kamchatka-Kuril-Japan-Izu-Mariana-Yap 140.1979 34.5512 190 22 kisz-32b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141.0551 33.0921 180 32 kisz-33b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141.0924 32.1047 173.8 27.65 kisz-33b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141.0924 32.1047 173.8 27.65 kisz-34a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141.6585 31.2408 172.1 25 kisz-34b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141.4154 30.1707 163 25 kisz-35a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141.4662 30.2899 163 1403 kisz-36a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.0670 29.4012 161.7 25.73 kisz-36b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.0670 29.4012 161.7 15.91 kisz-37a Kamchatka-Kuril-Japan-Izu-Mariana-Yap <t< td=""><td>97.02</td></t<>	97.02
kisz-31z Kamchatka-Kuril-Japan-Izu-Mariana-Yap 140.1979 34.5512 190 22 kisz-32a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141.0551 33.0921 180 32 kisz-32b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141.5098 33.0921 180 21.69 kisz-33b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141.5098 32.1047 173.8 27.65 kisz-34b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141.6596 32.1473 173.8 18.27 kisz-34b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141.6585 31.2408 172.1 25 kisz-35a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141.6585 31.2408 172.1 15.38 kisz-35b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141.6261 30.2899 163 14.03 kisz-36b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.0670 29.4012 161.7 25.73 kisz-37b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.2067 29.4012 161.7 15.91 kisz-38b Kamchatka-Kuril-Japan-Izu-Mariana-Yap	78.29
kisz-32a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141.0551 33.0921 180 32 kisz-32b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141.0924 32.1047 173.8 27.65 kisz-33b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141.0924 32.1047 173.8 27.65 kisz-34b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141.5596 32.1473 173.8 18.27 kisz-34a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141.16585 31.2408 172.1 15.38 kisz-34b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141.15689 31.1851 172.1 25 kisz-35a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141.6662 30.2899 163 14.03 kisz-35a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141.6261 29.2740 161.7 25.73 kisz-36a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.0670 29.4012 161.7 25.73 kisz-37a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.20670 29.4012 161.7 15.91 kisz-38a Kamchatka-Kuril-Japan-Izu-Mariana-	59.56
kisz-32b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141.5098 33.0921 180 21.69 kisz-33a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141.0924 32.1047 173.8 27.65 kisz-33b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141.5596 32.1473 173.8 1827 kisz-34b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141.1689 31.1851 172.1 25 kisz-35a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141.6585 31.2408 172.1 15.38 kisz-35a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141.6662 30.2899 163 25 kisz-35a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141.6261 29.2740 161.7 25.73 kisz-36a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.0670 29.4012 161.7 15.91 kisz-36b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.0670 29.4012 161.7 15.91 kisz-37a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.2254 27.6946 170.3 20 kisz-38b Kamchatka-Kuril-Japan-Izu-Mariana-Yap <td>40.83</td>	40.83
kisz-33a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141.0924 32.1047 173.8 27.65 kisz-34b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141.5596 32.1473 173.8 18.27 kisz-34b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141.1869 31.1851 172.1 25 kisz-35a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141.6685 31.2408 172.1 15.38 kisz-35b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141.8662 30.2899 163 14.03 kisz-36a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.0670 29.4012 161.7 25.73 kisz-36a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.0120 29.3322 154.7 20 kisz-36b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.0120 28.3322 154.7 20 kisz-37b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.0120 28.3322 154.7 20 kisz-38a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.0120 28.6124 17.70 11 kisz-39a Kamchatka-Kuril-Japan-Izu-Mariana-Yap	23.48
kisz-33b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141.5596 32.1473 173.8 18.27 kisz-34a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141.1869 31.1851 172.1 25 kisz-34b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141.6585 31.2408 172.1 15.38 kisz-35b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141.6662 30.2899 163 14.03 kisz-36a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141.6261 29.2740 161.7 25.73 kisz-36a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.0670 29.4012 161.7 25.73 kisz-37a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.0120 28.3322 154.7 20 kisz-38a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.2653 27.6946 170.3 20 kisz-39a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.26955 27.7659 170.3 11 kisz-39a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.6673 26.9127 177.2 24.23 kisz-40a Kamchatka-Kuril-Japan-Izu-Mariana-Yap <td>5</td>	5
kisz-34a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141,1869 31,1851 172,1 25 kisz-34b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141,6585 31,2408 172,1 15,38 kisz-35a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141,6585 31,2408 172,1 15,38 kisz-35b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141,6562 30,2899 163 140,33 kisz-36b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141,6261 29,2740 161,7 25,73 kisz-36b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142,0670 29,4012 161,7 15,91 kisz-37a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142,0120 28,3322 154,7 20 kisz-37a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142,6070 29,4012 161,7 15,91 kisz-37a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142,0120 28,3322 154,7 11 kisz-38a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142,6073 26,9127 177,2 24,23 kisz-39b Kamchatka-Kuril-Japan-Izu-Mariana-Yap	20.67
kisz-34b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141.6585 31.2408 172.1 15.38 kisz-35a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141.4154 30.1707 163 25 kisz-35b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141.6661 30.2899 163 14.03 kisz-36b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141.6661 29.2740 161.7 25.73 kisz-37b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.0670 29.4012 161.7 25.73 kisz-37b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.0120 28.3322 154.7 20 kisz-38b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.2254 27.6946 170.3 20 kisz-38b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.26955 27.7659 170.3 11 kisz-39a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.7674 26.9325 177.2 24.23 kisz-40a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.7674 26.9325 177.2 14.38 kisz-41b Kamchatka-Kuril-Japan-Izu-Mariana-Yap	5
kisz-35a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141.4154 30.1707 163 25 kisz-35b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141.8662 30.2899 163 14.03 kisz-36a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141.6261 29.2740 161.7 25.73 kisz-37b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.0670 29.4012 161.7 15.91 kisz-37b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.0120 28.3322 154.7 20 kisz-38a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.2254 27.6946 170.3 20 kisz-38b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.6955 27.7659 170.3 11 kisz-39b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.7674 26.9325 177.2 24.23 kisz-40a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.2673 26.1923 189.4 26.49 kisz-41b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.7090 26.1264 189.4 20.2 kisz-42b Kamchatka-Kuril-Japan-Izu-Mariana-Yap	18.26
kisz–35b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141.8662 30.2899 163 14.03 kisz–36a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141.6261 29.2740 161.7 25.73 kisz–36b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.0670 29.4012 161.7 15.91 kisz–37a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.0120 28.3322 154.7 20 kisz–37b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.0202 28.5124 154.7 11 kisz–38a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.2254 27.6946 170.3 20 kisz–39a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.6955 27.7659 170.3 11 kisz–39b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.2673 26.9127 177.2 24.23 kisz–40a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.7674 26.9325 177.2 14.38 kisz–40b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.6165 25.1184 173.7 22.07 kisz–41b Kamchatka-Kuril-Japan-Izu-Mariana-Yap <td>5</td>	5
kisz–36a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 141.6261 29.2740 161.7 25.73 kisz–36b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.0670 29.4012 161.7 15.91 kisz–37a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.0120 28.3322 154.7 20 kisz–37b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.4463 28.5124 154.7 11 kisz–38a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.2254 27.6696 170.3 20 kisz–39a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.6955 27.7659 170.3 11 kisz–39b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.7674 26.9325 177.2 24.23 kisz–40a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.2673 26.1923 189.4 26.49 kisz–40b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.27090 26.1264 189.4 20.2 kisz–41b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.6165 25.1184 173.7 16.36 kisz–42a Kamchatka-Kuril-Japan-Izu-Mariana-Yap </td <td>17.12</td>	17.12
kisz–36b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.0670 29.4012 161.7 15.91 kisz–37a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.0120 28.3322 154.7 20 kisz–37b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.0120 28.3322 154.7 20 kisz–37b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.0463 28.5124 154.7 11 kisz–38a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.2254 27.6946 170.3 20 kisz–39b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.6955 27.7659 170.3 11 kisz–39b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.2673 26.9127 177.2 24.23 kisz–40a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.2673 26.1923 189.4 26.49 kisz–41b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.7090 26.1264 189.4 20.2 kisz–42a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.6165 25.1184 173.7 16.36 kisz–43b Kamchatka-Kuril-Japan-Izu-Mariana-Yap	5
kisz-37a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.0120 28.3322 154.7 20 kisz-37b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.4463 28.5124 154.7 11 kisz-38a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.2254 27.6946 170.3 20 kisz-38b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.6955 27.7659 170.3 11 kisz-39a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.3085 26.9127 177.2 24.23 kisz-39b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.7674 26.9325 177.2 24.23 kisz-40a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.7674 26.9325 177.2 24.23 kisz-40b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.7090 26.1264 189.4 20.2 kisz-41b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.6165 25.1184 173.7 16.36 kisz-42b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 143.1321 24.1432 143.5 21.54 kisz-43b Kamchatka-Kuril-Japan-Izu-Mariana-Yap <td>18.71</td>	18.71
kisz–37b Kamchatka-Kuril-Japan-Izu-Mariana-Yap kisz–38a Kamchatka-Kuril-Japan-Izu-Mariana-Yap kisz–38b Kamchatka-Kuril-Japan-Izu-Mariana-Yap kisz–39b Kamchatka-Kuril-Japan-Izu-Mariana-Yap kisz–39a Kamchatka-Kuril-Japan-Izu-Mariana-Yap kisz–39b Kamchatka-Kuril-Japan-Izu-Mariana-Yap kisz–39b Kamchatka-Kuril-Japan-Izu-Mariana-Yap kisz–40a Kamchatka-Kuril-Japan-Izu-Mariana-Yap kisz–40b Kamchatka-Kuril-Japan-Izu-Mariana-Yap kisz–41a Kamchatka-Kuril-Japan-Izu-Mariana-Yap kisz–41b Kamchatka-Kuril-Japan-Izu-Mariana-Yap kisz–42b Kamchatka-Kuril-Japan-Izu-Mariana-Yap kisz–42b Kamchatka-Kuril-Japan-Izu-Mariana-Yap kisz–42b Kamchatka-Kuril-Japan-Izu-Mariana-Yap kisz–43a Kamchatka-Kuril-Japan-Izu-Mariana-Yap kisz–43b Kamchatka-Kuril-Japan-Izu-Mariana-Yap kisz–44b Kamchatka-Kuril-Japan-Izu-Mariana-Yap kisz–44b Kamchatka-Kuril-Japan-Izu-Mariana-Yap kisz–44b Kamchatka-Kuril-Japan-Izu-Mariana-Yap kisz–45b Kamchatka-Kuril-Japan-Izu-Mariana-Yap kisz–45b Kamchatka-Kuril-Japan-Izu-Mariana-Yap kisz–45b Kamchatka-Kuril-Japan-Izu-Mariana-Yap kisz–45b Kamchatka-Kuril-Japan-Izu-Mariana-Yap kisz–46a Kamchatka-Kuril-Japan-Izu-Mariana-Yap kisz–46b Kamchatka-Kuril-Japan-Izu-Mariana-Yap kisz–46b Kamchatka-Kuril-Japan-Izu-Mariana-Yap kisz–46b Kamchatka-Kuril-Japan-Izu-Mariana-Yap kisz–46b Kamchatka-Kuril-Japan-Izu-Mariana-Yap kisz–47b Kamchatka-Kuril-Japan-Izu-Mariana-Yap kisz–47b Kamchatka-Kuril-Japan-Izu-Mariana-Yap kisz–47b Kamchatka-Kuril-Japan-Izu-Mariana-Yap kisz–47b Kamchatka-Kuril-Japan-Izu-Mariana-Yap kisz–47b Kamchatka-Kuril-Japan-Izu-Mariana-Yap kisz–47b Kamchatka-Kuril-Japan-Izu-Mariana-Yap kisz–48b Kamchatka-Kuril-Japan-Izu-Mariana-Yap kisz–47b Kamchatka-Kuril-Japan-Izu-Mariana-Yap kisz–48b Kamchatka-Kuril-	5
kisz–38a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.2254 27.6946 170.3 20 kisz–38b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.6955 27.7659 170.3 11 kisz–39a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.3085 26.9127 177.2 24.23 kisz–39b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.7674 26.9325 177.2 14.38 kisz–40a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.7674 26.9325 177.2 14.38 kisz–40b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.7674 26.9325 177.2 14.38 kisz–40b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.7090 26.1264 189.4 20.2 kisz–41a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.1595 25.0729 173.7 22.07 kisz–41b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.6165 25.1184 173.7 16.36 kisz–42a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.7641 23.8947 143.5 21.54 kisz–42b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 143.1321 24.1432 143.5 15.54 kisz–43a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 143.5281 23.0423 129.2 23.02 kisz–43b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 143.8128 23.3626 129.2 15.99 kisz–44a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 144.2230 22.5240 134.6 28.24 kisz–44b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 145.0895 21.8866 125.8 36.73 kisz–45b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 145.0895 21.8866 125.8 20.84 kisz–46a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 145.6972 21.3783 135.9 30.75 kisz–46b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 145.9954 21.6469 135.9 18.22 kisz–47a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 145.9954 21.6469 135.9 18.22 kisz–47a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 146.0406 20.9341 160.1 29.87 kisz–47b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 146.0406 20.9341 160.1 29.87 kisz–48b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 146.0406 20.900 158 32.75 kisz–48b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 146.0406 20.900 158 32.75 kisz–48	14.54
kisz–38b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.6955 27.7659 170.3 11 kisz–39a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.3085 26.9127 177.2 24.23 kisz–39b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.7674 26.9325 177.2 14.38 kisz–40a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.2673 26.1923 189.4 26.49 kisz–40b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.7090 26.1264 189.4 20.2 kisz–41a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.1595 25.0729 173.7 22.07 kisz–41b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.6165 25.1184 173.7 16.36 kisz–42a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.7641 23.8947 143.5 21.54 kisz–42b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 143.1321 24.1432 143.5 15.54 kisz–43a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 143.5281 23.0423 129.2 23.02 kisz–44b Kamchatka-Kuril-Japan-Izu-Maria	5
kisz–39a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.3085 26.9127 177.2 24.23 kisz–39b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.7674 26.9325 177.2 14.38 kisz–40a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.7674 26.9325 177.2 14.38 kisz–40b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.7090 26.1264 189.4 20.2 kisz–41a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.1595 25.0729 173.7 22.07 kisz–41b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.6165 25.1184 173.7 16.36 kisz–42a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.7641 23.8947 143.5 21.54 kisz–42b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 143.1321 24.1432 143.5 15.54 kisz–43a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 143.8128 23.3626 129.2 23.02 kisz–44b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 144.2230 22.5240 134.6 28.24 kisz–45b Kamchatka-Kuril-Japan-Izu-Ma	14.54
kisz–39b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.7674 26.9325 177.2 14.38 kisz–40a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.2673 26.1923 189.4 26.49 kisz–40b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.7090 26.1264 189.4 20.2 kisz–41a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.1595 25.0729 173.7 22.07 kisz–41b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.6165 25.1184 173.7 16.36 kisz–42a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.7641 23.8947 143.5 21.54 kisz–42b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 143.1321 24.1432 143.5 15.54 kisz–43b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 143.5281 23.0423 129.2 23.02 kisz–43b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 144.2230 22.5240 134.6 28.24 kisz–44b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 145.0895 21.8666 125.8 36.73 kisz–45b Kamchatka-Kuril-Japan-Izu-Ma	5
kisz–40a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.2673 26.1923 189.4 26.49 kisz–40b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.7090 26.1264 189.4 20.2 kisz–41a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.1595 25.0729 173.7 22.07 kisz–41b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.6165 25.1184 173.7 16.36 kisz–42a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.7641 23.8947 143.5 21.54 kisz–42b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 143.1321 24.1432 143.5 15.54 kisz–43a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 143.8128 23.3626 129.2 23.02 kisz–43b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 144.2230 22.5240 134.6 28.24 kisz–44b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 145.0895 21.8866 125.8 36.73 kisz–45a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 145.6972 21.3783 135.9 30.75 kisz–46a Kamchatka-Kuril-Japan-Izu-Ma	17.42
kisz–40b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.7090 26.1264 189.4 20.2 kisz–41a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.1595 25.0729 173.7 22.07 kisz–41b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.6165 25.1184 173.7 16.36 kisz–42a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.7641 23.8947 143.5 21.54 kisz–42b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 143.1321 24.1432 143.5 15.54 kisz–43a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 143.5281 23.0423 129.2 23.02 kisz–43b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 143.8128 23.3626 129.2 15.99 kisz–44a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 144.2230 22.5240 134.6 28.24 kisz–45a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 145.0895 21.8866 125.8 36.73 kisz–45b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 145.6972 21.3785 135.9 30.75 kisz–46b Kamchatka-Kuril-Japan-Izu-Ma	5
kisz–41a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.1595 25.0729 173.7 22.07 kisz–41b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.6165 25.1184 173.7 16.36 kisz–42a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.7641 23.8947 143.5 21.54 kisz–42b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 143.1321 24.1432 143.5 15.54 kisz–43a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 143.5281 23.0423 129.2 23.02 kisz–43b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 144.8128 23.3626 129.2 15.99 kisz–44a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 144.2230 22.5240 134.6 28.24 kisz–44b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 145.5246 22.8056 134.6 15.74 kisz–45a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 145.0895 21.8866 125.8 36.73 kisz–45b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 145.6972 21.3783 135.9 30.75 kisz–46b Kamchatka-Kuril-Japan-Izu-M	22.26 5
kisz–41b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.6165 25.1184 173.7 16.36 kisz–42a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.7641 23.8947 143.5 21.54 kisz–42b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 143.1321 24.1432 143.5 15.54 kisz–43a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 143.5281 23.0423 129.2 23.02 kisz–43b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 143.8128 23.3626 129.2 15.99 kisz–44b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 144.2230 22.5240 134.6 28.24 kisz–45a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 145.0895 21.8866 125.8 36.73 kisz–45b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 145.3171 22.1785 125.8 20.84 kisz–46a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 145.6972 21.3783 135.9 30.75 kisz–47b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 146.0406 20.9341 160.1 29.87 kisz–48a Kamchatka-Kuril-Japan-Izu-M	19.08
kisz–42a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 142.7641 23.8947 143.5 21.54 kisz–42b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 143.1321 24.1432 143.5 15.54 kisz–43a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 143.5281 23.0423 129.2 23.02 kisz–43b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 143.8128 23.3626 129.2 15.99 kisz–44a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 144.2230 22.5240 134.6 28.24 kisz–45a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 145.0895 21.8866 125.8 36.73 kisz–45b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 145.6972 21.3783 135.9 30.75 kisz–46a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 145.6972 21.3783 135.9 30.75 kisz–46b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 146.0406 20.9341 160.1 29.87 kisz–47a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 146.4330 21.0669 160.1 17 kisz–48a Kamchatka-Kuril-Japan-Izu-Mari	5
kisz–42b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 143.1321 24.1432 143.5 15.54 kisz–43a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 143.5281 23.0423 129.2 23.02 kisz–43b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 143.8128 23.3626 129.2 15.99 kisz–44a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 144.2230 22.5240 134.6 28.24 kisz–44b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 144.5246 22.8056 134.6 15.74 kisz–45a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 145.0895 21.8866 125.8 36.73 kisz–45b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 145.6972 21.3783 135.9 30.75 kisz–46b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 145.9954 21.6469 135.9 18.22 kisz–47a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 146.0406 20.9341 160.1 29.87 kisz–47b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 146.4330 21.0669 160.1 17 kisz–48a Kamchatka-Kuril-Japan-Izu-Mari	18.4
kisz–43a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 143.5281 23.0423 129.2 23.02 kisz–43b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 143.8128 23.3626 129.2 15.99 kisz–44a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 144.2230 22.5240 134.6 28.24 kisz–44b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 144.5246 22.8056 134.6 15.74 kisz–45a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 145.0895 21.8866 125.8 36.73 kisz–45b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 145.3171 22.1785 125.8 20.84 kisz–46a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 145.6972 21.3783 135.9 30.75 kisz–46b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 145.9954 21.6469 135.9 18.22 kisz–47a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 146.0406 20.9341 160.1 29.87 kisz–47b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 146.330 21.0669 160.1 17 kisz–48a Kamchatka-Kuril-Japan-Izu-Maria	5
kisz-43b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 143.8128 23.3626 129.2 15.99 kisz-44a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 144.2230 22.5240 134.6 28.24 kisz-44b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 144.5246 22.8056 134.6 15.74 kisz-45a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 145.0895 21.8866 125.8 36.73 kisz-45b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 145.3171 22.1785 125.8 20.84 kisz-46a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 145.6972 21.3783 135.9 30.75 kisz-46b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 146.0406 20.9341 160.1 29.87 kisz-47b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 146.4330 21.0669 160.1 17 kisz-48a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 146.3836 20.0690 158 32.75 kisz-48b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 146.7567 20.2108 158 17.07	18.77
kisz-44a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 144.2230 22.5240 134.6 28.24 kisz-44b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 144.5246 22.8056 134.6 15.74 kisz-45a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 145.0895 21.8866 125.8 36.73 kisz-45b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 145.3171 22.1785 125.8 20.84 kisz-46a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 145.6972 21.3783 135.9 30.75 kisz-46b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 146.0406 20.9341 160.1 29.87 kisz-47b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 146.4330 21.0669 160.1 17 kisz-48a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 146.3836 20.0690 158 32.75 kisz-48b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 146.7567 20.2108 158 17.07	5
kisz-44b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 144.5246 22.8056 134.6 15.74 kisz-45a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 145.0895 21.8866 125.8 36.73 kisz-45b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 145.3171 22.1785 125.8 20.84 kisz-46a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 145.6972 21.3783 135.9 30.75 kisz-46b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 145.9954 21.6469 135.9 18.22 kisz-47a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 146.0406 20.9341 160.1 29.87 kisz-47b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 146.330 21.0669 160.1 17 kisz-48a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 146.3836 20.0690 158 32.75 kisz-48b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 146.7567 20.2108 158 17.07	18.56
kisz–45a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 145.0895 21.8866 125.8 36.73 kisz–45b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 145.3171 22.1785 125.8 20.84 kisz–46a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 145.6972 21.3783 135.9 30.75 kisz–46b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 145.9954 21.6469 135.9 18.22 kisz–47a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 146.0406 20.9341 160.1 29.87 kisz–47b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 146.330 21.0669 160.1 17 kisz–48a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 146.3836 20.0690 158 32.75 kisz–48b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 146.7567 20.2108 158 17.07	5
kisz–45b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 145.3171 22.1785 125.8 20.84 kisz–46a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 145.6972 21.3783 135.9 30.75 kisz–46b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 145.9954 21.6469 135.9 18.22 kisz–47a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 146.0406 20.9341 160.1 29.87 kisz–47b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 146.330 21.0669 160.1 17 kisz–48a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 146.3836 20.0690 158 32.75 kisz–48b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 146.7567 20.2108 158 17.07	22.79
kisz–46b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 145.9954 21.6469 135.9 18.22 kisz–47a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 146.0406 20.9341 160.1 29.87 kisz–47b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 146.4330 21.0669 160.1 17 kisz–48a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 146.3836 20.0690 158 32.75 kisz–48b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 146.7567 20.2108 158 17.07	5
kisz–46b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 145.9954 21.6469 135.9 18.22 kisz–47a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 146.0406 20.9341 160.1 29.87 kisz–47b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 146.4330 21.0669 160.1 17 kisz–48a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 146.3836 20.0690 158 32.75 kisz–48b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 146.7567 20.2108 158 17.07	20.63
kisz–47b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 146.4330 21.0669 160.1 17 kisz–48a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 146.3836 20.0690 158 32.75 kisz–48b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 146.7567 20.2108 158 17.07	5
kisz–48a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 146.3836 20.0690 158 32.75 kisz–48b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 146.7567 20.2108 158 17.07	19.62
kisz–48b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 146.7567 20.2108 158 17.07	5
	19.68
kieg 40a Vemehatka Vuril Ianan Igu Mariana Van 146 0000 10 2122 104 5 25 07	5
kisz–49a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 146.6689 19.3123 164.5 25.07	21.41
kisz–49b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 147.0846 19.4212 164.5 19.16	5
kisz–50a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 146.9297 18.5663 172.1 22	22.1
kisz–50b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 147.3650 18.6238 172.1 20	5
kisz–51a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 146.9495 17.7148 175.1 22.06	22.04
kisz–51b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 147.3850 17.7503 175.1 19.93	5
kisz–52a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 146.9447 16.8869 180 25.51	18.61
kisz–52b Kamchatka-Kuril-Japan-Izu-Mariana-Yap 147.3683 16.8869 180 15.79	5
kisz–53a Kamchatka-Kuril-Japan-Izu-Mariana-Yap 146.8626 16.0669 185.2 27.39	18.41
Continued on	next page

Table B.5 – continued

Segment	Description	Longitude(°E)	Latitude(^o N)	Strike(0)	Dip(°)	Depth (km)
kisz–53b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	147.2758	16.0309	185.2	15.56	5
kisz-54a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap		15.3883	199.1	28.12	20.91
kisz-54b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap		15.2590	199.1	18.56	5
kisz-55a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	146.4717	14.6025	204.3	29.6	26.27
kisz-55b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	146.8391	14.4415	204.3	25.18	5
kisz-56a	Kamchatka-Kuril-Japan-Izu-Mariana-Yar	146.1678	13.9485	217.4	32.04	26.79
kisz-56b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	146.4789	13.7170	217.4	25.84	5
kisz-57a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	145.6515	13.5576	235.8	37	24.54
kisz-57b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	145.8586	13.2609	235.8	23	5
kisz–58a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	144.9648	12.9990	237.8	37.72	24.54
kisz–58b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	145.1589	12.6984	237.8	23	5
kisz-59a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	144.1799	12.6914	242.9	34.33	22.31
kisz–59b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	144.3531	12.3613	242.9	20.25	5
kisz-60a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	143.3687	12.3280	244.9	30.9	20.62
kisz-60b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	143.5355	11.9788	244.9	18.2	5
kisz–61a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	142.7051	12.1507	261.8	35.41	25.51
kisz–61b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	142.7582	11.7883	261.8	24.22	5
kisz-62a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	141.6301	11.8447	245.7	39.86	34.35
kisz-62b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	141.7750	11.5305	245.7	35.94	5
kisz-63a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	140.8923	11.5740	256.2	42	38.46
kisz-63b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	140.9735	11.2498	256.2	42	5
kisz-64a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	140.1387	11.6028	269.6	42.48	38.77
kisz-64b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	140.1410	11.2716	269.6	42.48	5
kisz-65a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	139.4595	11.5883	288.7	44.16	39.83
kisz–65b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	139.3541	11.2831	288.7	44.16	5
kisz–66a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	138.1823	11.2648	193.1	45	40.36
kisz–66b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	138.4977	11.1929	193.1	45	5
kisz-67a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap		10.3398	189.8	45	40.36
kisz–67b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap		10.2856	189.8	45	5
kisz–68a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap		9.6136	201.7	45	40.36
kisz–68b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	138.0599	9.4963	201.7	45	5
kisz–69a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap		8.8996	213.5	45	40.36
kisz–69b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap		8.7241	213.5	45	5
kisz–70a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap		8.2872	226.5	45	40.36
kisz–70b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	137.2400	8.0569	226.5	45	5
kisz–71a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap		7.9078	263.9	45	40.36
kisz–71b	Kamchatka-Kuril-Japan-Izu-Mariana-Yar	136.4202	7.5920	263.9	45	5
kisz–72a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap		7.9130	276.9	45	40.36
kisz–72b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap		7.5977	276.9	45	5
kisz-73a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap		7.4541	224	45	40.36
kisz–73b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap		7.2335	224	45	5
kisz–74a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap		6.8621	228.1	45	40.36
kisz–74b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap		6.6258	228.1	45	5
kisz–75a	Kamchatka-Kuril-Japan-Izu-Mariana-Yap		6.1221	217.7	45	40.36
kisz–75b	Kamchatka-Kuril-Japan-Izu-Mariana-Yap	133.2751	5.9280	217.7	45	5



Figure B.6: Manus–Oceanic Convergent Boundary unit sources.

Table B.6: Earthquake parameters for Manus–Oceanic Convergent Boundary unit sources.

Segment	Description	Longitude(^o E)	Latitude(^o N)	Strike(0)	Dip(o)	Depth (km)
mosz–1a	Manus-Oceanic Convergent Boundary	154.0737	-4.8960	140.2	15	15.88
mosz–1b	Manus-Oceanic Convergent Boundary	154.4082	-4.6185	140.2	15	2.94
mosz–2a	Manus-Oceanic Convergent Boundary	153.5589	-4.1575	140.2	15	15.91
mosz–2b	Manus-Oceanic Convergent Boundary	153.8931	-3.8800	140.2	15	2.97
mosz–3a	Manus-Oceanic Convergent Boundary	153.0151	-3.3716	143.9	15	16.64
mosz–3b	Manus-Oceanic Convergent Boundary	153.3662	-3.1160	143.9	15	3.7
mosz–4a	Manus-Oceanic Convergent Boundary	152.4667	-3.0241	127.7	15	17.32
mosz–4b	Manus-Oceanic Convergent Boundary	152.7321	-2.6806	127.7	15	4.38
mosz–5a	Manus-Oceanic Convergent Boundary	151.8447	-2.7066	114.3	15	17.57
mosz–5b	Manus-Oceanic Convergent Boundary	152.0235	-2.3112	114.3	15	4.63
mosz–6a	Manus-Oceanic Convergent Boundary	151.0679	-2.2550	115	15	17.66
mosz–6b	Manus-Oceanic Convergent Boundary	151.2513	-1.8618	115	15	4.72
mosz–7a	Manus-Oceanic Convergent Boundary	150.3210	-2.0236	107.2	15	17.73
mosz–7b	Manus-Oceanic Convergent Boundary	150.4493	-1.6092	107.2	15	4.79
mosz–8a	Manus-Oceanic Convergent Boundary	149.3226	-1.6666	117.8	15	17.83
mosz–8b	Manus-Oceanic Convergent Boundary	149.5251	-1.2829	117.8	15	4.89
mosz–9a	Manus-Oceanic Convergent Boundary	148.5865	-1.3017	112.7	15	17.84
mosz–9b	Manus-Oceanic Convergent Boundary	148.7540	-0.9015	112.7	15	4.9
mosz–10a	Manus-Oceanic Convergent Boundary	147.7760	-1.1560	108	15	17.78
mosz–10b	Manus-Oceanic Convergent Boundary	147.9102	-0.7434	108	15	4.84
mosz–11a	Manus-Oceanic Convergent Boundary	146.9596	-1.1226	102.5	15	17.54
mosz–11b	Manus-Oceanic Convergent Boundary	147.0531	-0.6990	102.5	15	4.6
mosz–12a	Manus-Oceanic Convergent Boundary	146.2858	-1.1820	87.48	15	17.29
mosz–12b	Manus-Oceanic Convergent Boundary	146.2667	-0.7486	87.48	15	4.35
mosz–13a	Manus-Oceanic Convergent Boundary	145.4540	-1.3214	83.75	15	17.34
mosz–13b	Manus-Oceanic Convergent Boundary	145.4068	-0.8901	83.75	15	4.4
mosz–14a	Manus-Oceanic Convergent Boundary	144.7151	-1.5346	75.09	15	17.21
mosz–14b	Manus-Oceanic Convergent Boundary	144.6035	-1.1154	75.09	15	4.27
mosz–15a	Manus-Oceanic Convergent Boundary	143.9394	-1.8278	70.43	15	16.52
mosz–15b	Manus-Oceanic Convergent Boundary	143.7940	-1.4190	70.43	15	3.58
mosz–16a	Manus-Oceanic Convergent Boundary	143.4850	-2.2118	50.79	15	15.86
mosz–16b	Manus-Oceanic Convergent Boundary	143.2106	-1.8756	50.79	15	2.92
mosz–17a	Manus-Oceanic Convergent Boundary	143.1655	-2.7580	33	15	16.64
mosz–17b	Manus-Oceanic Convergent Boundary	142.8013	-2.5217	33	15	3.7




Figure B.7: New Guinea Subduction Zone unit sources.

Table B.7: Earthquake parameters for New Guinea Subduction Zone unit sources.

Segment	Description	Longitude(^o E)	Latitude(^o N)	Strike(°)	Dip(o)	Depth (km)
ngsz–1a	New Guinea	143.6063	-4.3804	120	29	25.64
ngsz–1b	New Guinea	143.8032	-4.0402	120	29	1.4
ngsz–2a	New Guinea	142.9310	-3.9263	114	27.63	20.1
ngsz–2b	New Guinea	143.0932	-3.5628	114	21.72	1.6
ngsz–3a	New Guinea	142.1076	-3.5632	114	20.06	18.73
ngsz–3b	New Guinea	142.2795	-3.1778	114	15.94	5
ngsz–4a	New Guinea	141.2681	-3.2376	114	21	17.76
ngsz–4b	New Guinea	141.4389	-2.8545	114	14.79	5
ngsz–5a	New Guinea	140.4592	-2.8429	114	21.26	16.14
ngsz–5b	New Guinea	140.6296	-2.4605	114	12.87	5
ngsz–6a	New Guinea	139.6288	-2.4960	114	22.72	15.4
ngsz–6b	New Guinea	139.7974	-2.1175	114	12	5
ngsz–7a	New Guinea	138.8074	-2.1312	114	21.39	15.4
ngsz–7b	New Guinea	138.9776	-1.7491	114	12	5
ngsz–8a	New Guinea	138.0185	-1.7353	113.1	18.79	15.14
ngsz–8b	New Guinea	138.1853	-1.3441	113.1	11.7	5
ngsz–9a	New Guinea	137.1805	-1.5037	111	15.24	13.23
ngsz–9b	New Guinea	137.3358	-1.0991	111	9.47	5
ngsz–10a	New Guinea	136.3418	-1.1774	111	13.51	11.09
ngsz–10b	New Guinea	136.4983	-0.7697	111	7	5
ngsz–11a	New Guinea	135.4984	-0.8641	111	11.38	12.49
ngsz–11b	New Guinea	135.6562	-0.4530	111	8.62	5
ngsz–12a	New Guinea	134.6759	-0.5216	110.5	10	13.68
ngsz–12b	New Guinea	134.8307	-0.1072	110.5	10	5
ngsz–13a	New Guinea	133.3065	-1.0298	99.5	10	13.68
ngsz–13b	New Guinea	133.3795	-0.5935	99.5	10	5
ngsz–14a	New Guinea	132.4048	-0.8816	99.5	10	13.68
ngsz–14b	New Guinea	132.4778	-0.4453	99.5	10	5
ngsz–15a	New Guinea	131.5141	-0.7353	99.5	10	13.68
ngsz–15b	New Guinea	131.5871	-0.2990	99.5	10	5




Figure B.8: New Zealand–Kermadec–Tonga Subduction Zone unit sources.

 ${\it Table~B.8: Earth quake~parameters~for~New~Zealand-Kermadec-Tonga~Subduction~Zone~unit~sources.}$

Segment	Description	Longitude(°E)	Latitude(^o N)	Strike(0)	Dip(°)	Depth (km)
ntsz-1a	New Zealand-Kermadec-Tonga	174.0985	-41.3951	258.6	24	25.34
ntsz–1b	New Zealand–Kermadec–Tonga	174.2076	-41.7973	258.6	24	5
ntsz–2a	New Zealand–Kermadec–Tonga	175.3289	-41.2592	260.6	29.38	23.17
ntsz–2b	New Zealand–Kermadec–Tonga	175.4142	-41.6454	260.6	21.31	5
ntsz–3a	New Zealand-Kermadec-Tonga	176.2855	-40.9950	250.7	29.54	21.74
ntsz–3b	New Zealand–Kermadec–Tonga	176.4580	-41.3637	250.7	19.56	5
ntsz–4a	New Zealand–Kermadec–Tonga	177.0023	-40.7679	229.4	24.43	18.87
ntsz–4b	New Zealand–Kermadec–Tonga	177.3552	-41.0785	229.4	16.1	5
ntsz–5a	New Zealand–Kermadec–Tonga	177.4114	-40.2396	210	18.8	19.29
ntsz–5b	New Zealand–Kermadec–Tonga	177.8951	-40.4525	210	16.61	5
ntsz–6a	New Zealand–Kermadec–Tonga	177.8036	-39.6085	196.7	18.17	15.8
ntsz–6b	New Zealand–Kermadec–Tonga	178.3352	-39.7310	196.7	12.48	5
ntsz–7a	New Zealand–Kermadec–Tonga	178.1676	-38.7480	197	28.1	17.85
ntsz–7b	New Zealand–Kermadec–Tonga	178.6541	-38.8640	197	14.89	5
ntsz–8a	New Zealand–Kermadec–Tonga	178.6263	-37.8501	201.4	31.47	18.78
ntsz–8b	New Zealand–Kermadec–Tonga	179.0788	-37.9899	201.4	16	5
ntsz–9a	New Zealand–Kermadec–Tonga	178.9833	-36.9770	202.2	29.58	20.02
ntsz–9b	New Zealand–Kermadec–Tonga	179.4369	-37.1245	202.2	17.48	5
ntsz–10a	New Zealand–Kermadec–Tonga	179.5534	-36.0655	210.6	32.1	20.72
ntsz–10b	New Zealand–Kermadec–Tonga	179.9595	-36.2593	210.6	18.32	5
ntsz–11a	New Zealand–Kermadec–Tonga	179.9267	-35.3538	201.7	25	16.09
ntsz–11b	New Zealand–Kermadec–Tonga	180.3915	-35.5040	201.7	12.81	5
ntsz–12a	New Zealand–Kermadec–Tonga	180.4433	-34.5759	201.2	25	15.46
ntsz–12b	New Zealand–Kermadec–Tonga	180.9051	-34.7230	201.2	12.08	5
ntsz–13a	New Zealand–Kermadec–Tonga	180.7990	-33.7707	199.8	25.87	19.06
ntsz–13b	New Zealand–Kermadec–Tonga	181.2573	-33.9073	199.8	16.33	5
ntsz–14a	New Zealand–Kermadec–Tonga	181.2828	-32.9288	202.4	31.28	22.73
ntsz–14b	New Zealand–Kermadec–Tonga	181.7063	-33.0751	202.4	20.77	5
ntsz–15a	New Zealand–Kermadec–Tonga	181.4918	-32.0035	205.4	32.33	22.64
ntsz–15b	New Zealand–Kermadec–Tonga	181.8967	-32.1665	205.4	20.66	5
ntsz–16a	New Zealand-Kermadec-Tonga	181.9781	-31.2535	205.5	34.29	23.59
ntsz–16b	New Zealand-Kermadec-Tonga	182.3706	-31.4131	205.5	21.83	5
ntsz–17a	New Zealand-Kermadec-Tonga	182.4819	-30.3859	210.3	37.6	25.58
ntsz–17b	New Zealand–Kermadec–Tonga	182.8387	-30.5655	210.3	24.3	5
ntsz–18a	New Zealand–Kermadec–Tonga	182.8176	-29.6545	201.6	37.65	26.13
ntsz–18b	New Zealand–Kermadec–Tonga	183.1985	-29.7856	201.6	25	5
ntsz–19a	New Zealand-Kermadec-Tonga	183.0622	-28.8739	195.7	34.41	26.13
ntsz–19b	New Zealand-Kermadec-Tonga	183.4700	-28.9742	195.7	25	5
ntsz–20a	New Zealand–Kermadec–Tonga	183.2724	-28.0967	188.8	38	26.13
ntsz–20b	New Zealand–Kermadec–Tonga	183.6691	-28.1508	188.8	25	5
ntsz–21a	New Zealand–Kermadec–Tonga	183.5747	-27.1402	197.1	32.29	24.83
ntsz–21b	New Zealand–Kermadec–Tonga	183.9829	-27.2518	197.1	23.37	5
ntsz–22a	New Zealand–Kermadec–Tonga	183.6608	-26.4975	180	29.56	18.63
ntsz–22b	New Zealand–Kermadec–Tonga	184.0974	-26.4975	180	15.82	5
ntsz–23a	New Zealand–Kermadec–Tonga	183.7599	-25.5371	185.8	32.42	20.56
ntsz–23b	New Zealand-Kermadec-Tonga	184.1781	-25.5752	185.8	18.13	5
ntsz–24a	New Zealand–Kermadec–Tonga	183.9139	-24.6201	188.2	33.31	23.73
ntsz–24b	New Zealand–Kermadec–Tonga	184.3228	-24.6734	188.2	22	5
ntsz–25a	New Zealand–Kermadec–Tonga	184.1266	-23.5922	198.5	29.34	19.64
ntsz–25b	New Zealand–Kermadec–Tonga	184.5322	-23.7163	198.5	17.03	5
ntsz–26a	New Zealand–Kermadec–Tonga	184.6613	-22.6460	211.7	30.26	19.43
ntsz–26b	New Zealand–Kermadec–Tonga	185.0196	-22.8497	211.7	16.78	5
ntsz–27a	New Zealand–Kermadec–Tonga	185.0879	-21.9139	207.9	31.73	20.67
ntsz–27b	New Zealand–Kermadec–Tonga	185.4522	-22.0928	207.9	18.27	5
ntsz–28a	New Zealand–Kermadec–Tonga	185.4037	-21.1758	200.5	32.44	21.76
ntsz–28b	New Zealand–Kermadec–Tonga	185.7849	-21.3084	200.5	19.58	5
ntsz–29a	New Zealand–Kermadec–Tonga	185.8087	-20.2629	206.4	32.47	20.4
ntsz-29b	New Zealand–Kermadec–Tonga	186.1710	-20.4312	206.4	17.94	5
ntsz–30a	New Zealand–Kermadec–Tonga	186.1499	-19.5087	200.9	32.98	22.46
ntsz-30b	New Zealand–Kermadec–Tonga	186.5236	-19.6432	200.9	20.44	5
				(Continued	on next page

Table B.8 – continued

Segment	Description	Longitude(^o E)	Latitude(^o N)	Strike(0)	Dip(°)	Depth (km)
ntsz-31a	New Zealand–Kermadec–Tonga	186.3538	-18.7332	193.9	34.41	21.19
ntsz-31b	New Zealand-Kermadec-Tonga	186.7339	-18.8221	193.9	18.89	5
ntsz–32a	New Zealand-Kermadec-Tonga	186.5949	-17.8587	194.1	30	19.12
ntsz-32b	New Zealand-Kermadec-Tonga	186.9914	-17.9536	194.1	16.4	5
ntsz-33a	New Zealand-Kermadec-Tonga	186.8172	-17.0581	190	33.15	23.34
ntsz-33b	New Zealand-Kermadec-Tonga	187.2047	-17.1237	190	21.52	5
ntsz–34a	New Zealand-Kermadec-Tonga	186.7814	-16.2598	182.1	15	13.41
ntsz-34b	New Zealand-Kermadec-Tonga	187.2330	-16.2759	182.1	9.68	5
ntsz-34c	New Zealand-Kermadec-Tonga	187.9697	-16.4956	7.62	57.06	6.571
ntsz–35a	New Zealand-Kermadec-Tonga	186.8000	-15.8563	149.8	15	12.17
ntsz–35b	New Zealand-Kermadec-Tonga	187.1896	-15.6384	149.8	8.24	5
ntsz-35c	New Zealand-Kermadec-Tonga	187.8776	-15.6325	342.4	57.06	6.571
ntsz–36a	New Zealand-Kermadec-Tonga	186.5406	-15.3862	123.9	40.44	36.72
ntsz-36b	New Zealand-Kermadec-Tonga	186.7381	-15.1025	123.9	39.38	5
ntsz-36c	New Zealand-Kermadec-Tonga	187.3791	-14.9234	307	57.06	6.571
ntsz–37a	New Zealand-Kermadec-Tonga	185.9883	-14.9861	102	68.94	30.99
ntsz–37b	New Zealand-Kermadec-Tonga	186.0229	-14.8282	102	31.32	5
ntsz–38a	New Zealand-Kermadec-Tonga	185.2067	-14.8259	88.4	80	26.13
ntsz-38b	New Zealand-Kermadec-Tonga	185.2044	-14.7479	88.4	25	5
ntsz–39a	New Zealand–Kermadec–Tonga	184.3412	-14.9409	82.55	80	26.13
ntsz-39b	New Zealand-Kermadec-Tonga	184.3307	-14.8636	82.55	25	5








Figure B.9: New Britain–Solomons–Vanuatu Zone unit sources.

 ${\it Table~B.9:~Earthquake~parameters~for~New~Britain-Solomons-Vanuatu~Subduction~Zone~unit~sources.}$

Segment	Description	Longitude(^o E)	Latitude(^o N)	Strike(o)	Dip(o)	Depth (km)
nvsz-1a	New Britain–Solomons–Vanuatu	148.6217	-6.4616	243.2	32.34	15.69
nvsz–1b	New Britain–Solomons–Vanuatu	148.7943	-6.8002	234.2	12.34	5
nvsz–2a	New Britain-Solomons-Vanuatu	149.7218	-6.1459	260.1	35.1	16.36
nvsz–2b	New Britain–Solomons–Vanuatu	149.7856	-6.5079	260.1	13.13	5
nvsz–3a	New Britain–Solomons–Vanuatu	150.4075	-5.9659	245.7	42.35	18.59
nvsz–3b	New Britain–Solomons–Vanuatu	150.5450	-6.2684	245.7	15.77	5
nvsz–4a	New Britain–Solomons–Vanuatu	151.1095	-5.5820	238.2	42.41	23.63
nvsz–4b	New Britain–Solomons–Vanuatu	151.2851	-5.8639	238.2	21.88	5
nvsz–5a	New Britain–Solomons–Vanuatu	152.0205	-5.1305	247.7	49.22	32.39
nvsz–5b	New Britain–Solomons–Vanuatu	152.1322	-5.4020	247.7	33.22	5
nvsz–6a	New Britain–Solomons–Vanuatu	153.3450	-5.1558	288.6	53.53	33.59
nvsz–6b	New Britain–Solomons–Vanuatu	153.2595	-5.4089	288.6	34.87	5
nvsz–7a	New Britain–Solomons–Vanuatu	154.3814	-5.6308	308.3	39.72	19.18
nvsz–7b	New Britain–Solomons–Vanuatu	154.1658	-5.9017	308.3	16.48	5
nvsz–8a	New Britain–Solomons–Vanuatu	155.1097	-6.3511	317.2	45.33	22.92
nvsz–8b	New Britain–Solomons–Vanuatu	154.8764	-6.5656	317.2	21	5
nvsz–9a	New Britain–Solomons–Vanuatu	155.5027	-6.7430	290.5	48.75	22.92
nvsz–9b	New Britain–Solomons–Vanuatu	155.3981	-7.0204	290.5	21	5
nvsz–10a	New Britain–Solomons–Vanuatu	156.4742	-7.2515	305.9	36.88	27.62
nvsz–10b	New Britain–Solomons–Vanuatu	156.2619	-7.5427	305.9	26.9	5
nvsz–11a	New Britain–Solomons–Vanuatu	157.0830	-7.8830	305.4	32.97	29.72
nvsz–11b	New Britain–Solomons–Vanuatu	156.8627	-8.1903	305.4	29.63	5
nvsz–12a	New Britain–Solomons–Vanuatu	157.6537	-8.1483	297.9	37.53	28.57
nvsz–12b	New Britain–Solomons–Vanuatu	157.4850	-8.4630	297.9	28.13	5
nvsz–13a	New Britain–Solomons–Vanuatu	158.5089	-8.5953	302.7	33.62	23.02
nvsz–13b	New Britain–Solomons–Vanuatu	158.3042	-8.9099	302.7	21.12	5
nvsz–14a	New Britain–Solomons–Vanuatu	159.1872	-8.9516	293.3	38.44	34.06
nvsz–14b	New Britain–Solomons–Vanuatu	159.0461	-9.2747	293.3	35.54	5
nvsz–15a	New Britain–Solomons–Vanuatu	159.9736	-9.5993	302.8	46.69	41.38
nvsz–15b	New Britain–Solomons–Vanuatu	159.8044	-9.8584	302.8	46.69	5
nvsz–16a	New Britain–Solomons–Vanuatu	160.7343	-10.0574	301	46.05	41
nvsz–16b	New Britain–Solomons–Vanuatu	160.5712	-10.3246	301	46.05	5
nvsz–17a	New Britain–Solomons–Vanuatu	161.4562	-10.5241	298.4	40.12	37.22
nvsz–17b	New Britain–Solomons–Vanuatu	161.2900	-10.8263	298.4	40.12	5
nvsz–18a	New Britain-Solomons-Vanuatu	162.0467	-10.6823	274.1	40.33	29.03
nvsz–18b	New Britain-Solomons-Vanuatu	162.0219	-11.0238	274.1	28.72	5
nvsz–19a	New Britain Solomons Vanuatu	162.7818	-10.5645	261.3	34.25	24.14
nvsz–19b nvsz–20a	New Britain–Solomons–Vanuatu New Britain–Solomons–Vanuatu	162.8392 163.7222	-10.9315	261.3 262.9	22.51	5 26.3
nvsz–20a nvsz–20b	New Britain–Solomons–Vanuatu		-10.5014 -10.7858	262.9	50.35 25.22	5
nvsz–200 nvsz–21a	New Britain–Solomons–Vanuatu	163.7581 164.9445	-10.7656	287.9	40.31	23.3
nvsz–21a	New Britain–Solomons–Vanuatu	164.8374	-10.7442	287.9	21.47	23.3 5
nvsz–210 nvsz–22a	New Britain–Solomons–Vanuatu	166.0261	-11.1069	317.1	42.39	20.78
nvsz–22b	New Britain–Solomons–Vanuatu	165.7783	-11.3328	317.1	18.4	5
nvsz–23a	New Britain–Solomons–Vanuatu	166.5179	-12.2260	342.4	47.95	22.43
nvsz–23a	New Britain–Solomons–Vanuatu	166.2244	-12.3171	342.4	20.4	5
nvsz–24a	New Britain–Solomons–Vanuatu	166.7236	-13.1065	342.4	47.13	28.52
nvsz–24a	New Britain–Solomons–Vanuatu	166.4241	-13.1979	342.6	28.06	5
nvsz–240 nvsz–25a	New Britain–Solomons–Vanuatu	166.8914	-13.1979	350.3	54.1	31.16
nvsz–25b	New Britain–Solomons–Vanuatu	166.6237	-14.0785 -14.1230	350.3	31.55	51.16
nvsz–250 nvsz–26a	New Britain–Solomons–Vanuatu	166.9200	-14.1250 -15.1450	365.6	50.46	29.05
nvsz–26b	New Britain–Solomons–Vanuatu	166.6252	-15.1170	365.6	28.75	29.03 5
nvsz–27a	New Britain–Solomons–Vanuatu	167.0053	-15.1170	334.2	44.74	25.46
nvsz–27a nvsz–27b	New Britain–Solomons–Vanuatu	166.7068	-15.6506 -15.7695	334.2	24.15	25.46 5
nvsz–27b nvsz–28a	New Britain–Solomons–Vanuatu	167.4074	-15.7695 -16.3455	327.5	41.53	22.44
nvsz–28b	New Britain–Solomons–Vanuatu	167.1117	-16.5455 -16.5264	327.5 327.5	20.42	5
nvsz–200 nvsz–29a	New Britain–Solomons–Vanuatu					3 24.12
	New Britain–Solomons–Vanuatu	167.9145 167.6229	-17.2807 -17.3757	341.2 341.2	49.1 22.48	24.12 5
nyez 20h		107.0229	-17.3737	341.4	44.40	
nvsz–29b nvsz–30a	New Britain–Solomons–Vanuatu	168.2220	-18.2353	348.6	44.19	23.99

Continued on next page

Table B.9 – continued

Segment	Description	Longitude(°E)	Latitude(^o N)	Strike(0)	Dip(o)	Depth (km)
nvsz–31a	New Britain–Solomons–Vanuatu	168.5022	-19.0510	345.6	42.2	22.26
nvsz-31b	New Britain-Solomons-Vanuatu	168.1611	-19.1338	345.6	20.2	5
nvsz-32a	New Britain-Solomons-Vanuatu	168.8775	-19.6724	331.1	42.03	21.68
nvsz-32b	New Britain-Solomons-Vanuatu	168.5671	-19.8338	331.1	19.49	5
nvsz-33a	New Britain-Solomons-Vanuatu	169.3422	-20.4892	332.9	40.25	22.4
nvsz-33b	New Britain-Solomons-Vanuatu	169.0161	-20.6453	332.9	20.37	5
nvsz-34a	New Britain-Solomons-Vanuatu	169.8304	-21.2121	329.1	39	22.73
nvsz-34b	New Britain-Solomons-Vanuatu	169.5086	-21.3911	329.1	20.77	5
nvsz-35a	New Britain-Solomons-Vanuatu	170.3119	-21.6945	311.9	39	22.13
nvsz–35b	New Britain-Solomons-Vanuatu	170.0606	-21.9543	311.9	20.03	5
nvsz-36a	New Britain-Solomons-Vanuatu	170.9487	-22.1585	300.4	39.42	23.5
nvsz-36b	New Britain-Solomons-Vanuatu	170.7585	-22.4577	300.4	21.71	5
nvsz-37a	New Britain-Solomons-Vanuatu	171.6335	-22.3087	281.3	30	22.1
nvsz–37b	New Britain–Solomons–Vanuatu	171.5512	-22.6902	281.3	20	5








Figure B.10: New Zealand–Puysegur Zone unit sources.

 $\label{lem:condition} \mbox{Table B.10: Earthquake parameters for New Zealand-Puysegur Subduction Zone unit sources.}$

Segment	Description	Longitude(^o E)	Latitude(^o N)	Strike(0)	Dip(o)	Depth (km)
nzsz-1a	New Zealand-Puysegur	168.0294	-45.4368	41.5	15	17.94
nzsz-1b	New Zealand-Puysegur	167.5675	-45.1493	41.5	15	5
nzsz-2a	New Zealand-Puysegur	167.3256	-46.0984	37.14	15	17.94
nzsz–2b	New Zealand-Puysegur	166.8280	-45.8365	37.14	15	5
nzsz-3a	New Zealand-Puysegur	166.4351	-46.7897	39.53	15	17.94
nzsz–3b	New Zealand-Puysegur	165.9476	-46.5136	39.53	15	5
nzsz–4a	New Zealand-Puysegur	166.0968	-47.2583	15.38	15	17.94
nzsz-4b	New Zealand-Puysegur	165.4810	-47.1432	15.38	15	5
nzsz–5a	New Zealand-Puysegur	165.7270	-48.0951	13.94	15	17.94
nzsz–5b	New Zealand-Puysegur	165.0971	-47.9906	13.94	15	5
nzsz-6a	New Zealand-Puysegur	165.3168	-49.0829	22.71	15	17.94
nzsz-6b	New Zealand-Puysegur	164.7067	-48.9154	22.71	15	5
nzsz-7a	New Zealand-Puysegur	164.8017	-49.9193	23.25	15	17.94
nzsz–7b	New Zealand-Puysegur	164.1836	-49.7480	23.25	15	5






Figure B.11: Ryukyu–Kyushu–Nankai Zone unit sources.

 ${\it Table~B.11:}~Earth quake~parameters~for~Ryukyu-Kyushu-Nankai~Subduction~Zone~unit~sources.$

Segment	Description	Longitude(^o E)	Latitude(^o N)	Strike(0)	Dip(o)	Depth (km)
rnsz-1a	Ryukyu–Kyushu–Nankai	122.6672	23.6696	262	14	11.88
rnsz-1b	Ryukyu–Kyushu–Nankai	122.7332	23.2380	262	10	3.2
rnsz-2a	Ryukyu-Kyushu-Nankai	123.5939	23.7929	259.9	18.11	12.28
rnsz–2b	Ryukyu–Kyushu–Nankai	123.6751	23.3725	259.9	10	3.6
rnsz–3a	Ryukyu–Kyushu–Nankai	124.4604	23.9777	254.6	19.27	14.65
rnsz–3b	Ryukyu–Kyushu–Nankai	124.5830	23.5689	254.6	12.18	4.1
rnsz–4a	Ryukyu-Kyushu-Nankai	125.2720	24.2102	246.8	18	20.38
rnsz–4b	Ryukyu–Kyushu–Nankai	125.4563	23.8177	246.8	16	6.6
rnsz–5a	Ryukyu–Kyushu–Nankai	125.9465	24.5085	233.6	18	20.21
rnsz–5b	Ryukyu–Kyushu–Nankai	126.2241	24.1645	233.6	16	6.43
rnsz–6a	Ryukyu-Kyushu-Nankai	126.6349	25.0402	228.7	17.16	19.55
rnsz-6b	Ryukyu-Kyushu-Nankai	126.9465	24.7176	228.7	15.16	6.47
rnsz–7a	Ryukyu-Kyushu-Nankai	127.2867	25.6343	224	15.85	17.98
rnsz–7b	Ryukyu–Kyushu–Nankai	127.6303	25.3339	224	13.56	6.26
rnsz–8a	Ryukyu-Kyushu-Nankai	128.0725	26.3146	229.7	14.55	14.31
rnsz–8b	Ryukyu-Kyushu-Nankai	128.3854	25.9831	229.7	9.64	5.94
rnsz–9a	Ryukyu-Kyushu-Nankai	128.6642	26.8177	219.2	15.4	12.62
rnsz–9b	Ryukyu–Kyushu–Nankai	129.0391	26.5438	219.2	8	5.66
rnsz-10a	Ryukyu–Kyushu–Nankai	129.2286	27.4879	215.2	17	12.55
rnsz-10b	Ryukyu–Kyushu–Nankai	129.6233	27.2402	215.2	8.16	5.45
rnsz–11a	Ryukyu-Kyushu-Nankai	129.6169	28.0741	201.3	17	12.91
rnsz–11b	Ryukyu-Kyushu-Nankai	130.0698	27.9181	201.3	8.8	5.26
rnsz–12a	Ryukyu–Kyushu–Nankai	130.6175	29.0900	236.7	16.42	13.05
rnsz-12b	Ryukyu–Kyushu–Nankai	130.8873	28.7299	236.7	9.57	4.74
rnsz–13a	Ryukyu–Kyushu–Nankai	130.7223	29.3465	195.2	20.25	15.89
rnsz-13b	Ryukyu-Kyushu-Nankai	131.1884	29.2362	195.2	12.98	4.66
rnsz–14a	Ryukyu-Kyushu-Nankai	131.3467	30.3899	215.1	22.16	19.73
rnsz–14b	Ryukyu-Kyushu-Nankai	131.7402	30.1507	215.1	17.48	4.71
rnsz–15a	Ryukyu-Kyushu-Nankai	131.9149	31.1450	216	15.11	16.12
rnsz–15b	Ryukyu-Kyushu-Nankai	132.3235	30.8899	216	13.46	4.48
rnsz–16a	Ryukyu-Kyushu-Nankai	132.5628	31.9468	220.9	10.81	10.88
rnsz–16b	Ryukyu-Kyushu-Nankai	132.9546	31.6579	220.9	7.19	4.62
rnsz–17a	Ryukyu-Kyushu-Nankai	133.6125	32.6956	239	10.14	12.01
rnsz–17b	Ryukyu-Kyushu-Nankai	133.8823	32.3168	239	8.41	4.7
rnsz–18a	Ryukyu-Kyushu-Nankai	134.6416	33.1488	244.7	10.99	14.21
rnsz-18b	Ryukyu-Kyushu-Nankai	134.8656	32.7502	244.5	10.97	4.7
rnsz–19a	Ryukyu-Kyushu-Nankai	135.6450	33.5008	246.5	14.49	14.72
rnsz-19b	Ryukyu-Kyushu-Nankai	135.8523	33.1021	246.5	11.87	4.44
rnsz-20a	Ryukyu-Kyushu-Nankai	136.5962	33.8506	244.8	15	14.38
rnsz–20b	Ryukyu–Kyushu–Nankai	136.8179	33.4581	244.8	12	3.98
rnsz–21a	Ryukyu-Kyushu-Nankai	137.2252	34.3094	231.9	15	15.4
rnsz–21b	Ryukyu-Kyushu-Nankai	137.5480	33.9680	231.9	12	5
rnsz–22a	Ryukyu-Kyushu-Nankai	137.4161	34.5249	192.3	15	15.4
rnsz–22b	Ryukyu-Kyushu-Nankai	137.9301	34.4327	192.3	12	5

Appendix C

SIFT Testing

C.1 Purpose

Forecast models are tested with synthetic tsunami events covering a range of tsunami source locations and magnitudes ranging from mega-events to microevents. Testing is also done with a selected set of historical tsunami events when available.

The purpose of forecast model testing is three-fold. The first objective is to assure that the results obtained with the Short-term Inundation Forecasting of Tsunamis (SIFT) software, which has been released to the Tsunami Warning Centers for operational use, agree with those obtained by the researcher during the development of the forecast model. The second objective is to test the forecast model for consistency, accuracy, time efficiency, and quality of results over a range of possible tsunami locations and magnitudes. The third objective is to identify bugs and issues in need of resolution by the researcher who developed the forecast model or by the SIFT software development team before the next version release to NOAA's two Tsunami Warning Centers.

Local hardware and software applications, and tools familiar to the researcher(s), are used to run the Method of Splitting Tsunami (MOST) model during the forecast model development. The test results presented in this report lend confidence that the model performs as developed and produces the same results when initiated within the SIFT application in an operational setting as those produced by the researcher(s) during the forecast model development. The test results assure those who rely on the Point Reyes tsunami forecast model that consistent results are produced irrespective of the system used.

C.2 Testing procedure

The general procedure for forecast model testing is to run a set of synthetic tsunami scenarios (and a selected set of historical tsunami events if available) through the SIFT application and compare the results with those obtained by the researcher during the forecast model development and presented in the tsunami forecast model report. Specific steps taken to test the model include:

• Identification of testing scenarios, including the standard set of synthetic events and customized synthetic scenarios that may have been used by the researcher(s) in developing the forecast model.

- Creation of new events to represent customized synthetic scenarios used by the researcher(s) in developing the forecast model, if any.
- Submission of test model runs with the forecast system, and export of the results from A, B, and C grids, along with time series.
- Recording applicable metadata, including the specific version of the forecast system used for testing.
- Examination of forecast model results from the forecast system for instabilities in both time series and plot results.
- Comparison of forecast model results obtained through the forecast system with those obtained during the forecast model development.
- Summarization of results with specific mention of quality, consistency, and time efficiency.
- Reporting of issues identified to modeler and forecast software development team.
- Retesting the forecast models in the forecast system when reported issues have been addressed or explained.

Synthetic model runs were tested on a DELL PowerEdge R510 computer equipped with two Xeon E5670 processors at 2.93 GHz, each with 12 MBytes of cache and 32GB memory. The processors are hex core and support hyperthreading, resulting in the computer performing as a 24 processor core machine. Additionally, the testing computer supports 10 Gigabit Ethernet for fast network connections. This computer configuration is similar or the same as the configurations of the computers installed at the Tsunami Warning Centers so the compute times should vary only slightly

C.3 Results

The Point Reyes, California forecast model was tested with five synthetic scenarios and one historical tsunami event. Test results from the forecast system and comparisons with the results obtained during the forecast model development are shown numerically in Table C.1 and graphically in Figures C.1 to C.6. The results show that the forecast model is stable and robust, with consistent and high quality results across geographically distributed tsunami sources and mega-event tsunami magnitudes. The model run time (wall clock time) was under 18 min for 8 hr of simulation time, and under 8 min for 4 hr thereby satisfying the "10 min run time per 4 hr of simulated time" criterion for operational efficiency.

Time series plots for two of the synthetic cases (CSSZ 89-98 and KISZ 22-31) were not present in the main report and their statistics were extracted from the original model output files from the development stage. The modeled scenarios were stable for all cases tested, with no instabilities or ringing. Results show that the largest modeled height was 401.25 cm and originated in the New Zealand-Kermadec-Tonga (NTSZ 30-39) source. Amplitudes greater than 100 cm were recorded for all synthetic test sources. The smallest signal of 119.8 cm was recorded for the far-field Central and South American (CSSZ 89-98) source. Direct comparisons, of output from the forecast tool with results of the historical event (Tohoku, referred to

as Honshu-2011 in the report) and available development synthetic events, demonstrated that the wave pattern were similar in shape, pattern and amplitude.






Figure C.1: Response of the Point Reyes forecast model to synthetic scenario KISZ 1-10 (α = 25). Maximum sea surface elevation for (a) A grid, (b) B grid, (c) C grid. Sea surface elevation time series at the C grid warning point (d). The extrema at the reference point are compared with the equivalent values obtained during model development in Figure 12.




Figure C.2: Response of the Point Reyes forecast model to synthetic scenario KISZ 22-31 (α = 25). Maximum sea surface elevation for (a) A grid, (b) B grid, (c) C grid. Sea surface elevation time series at the C grid warning point (d). The extrema at the reference point are compared with the equivalent values obtained during model development listed in Table C.1.




Figure C.3: Response of the Point Reyes forecast model to synthetic scenario ACSZ 56-65 (α =25). Maximum sea surface elevation for (a) A grid, (b) B grid, (c) C grid. Sea surface elevation time series at the C grid warning point (d). Panels (c) and (d) can be compared with the equivalent results, obtained during model development, displayed in Figure 11.




Figure C.4: Response of the Point Reyes forecast model to synthetic scenario CSSZ 89-98 (α = 25). Maximum sea surface elevation for (a) A grid, (b) B grid, (c) C grid. Sea surface elevation time series at the C grid warning point (d). Panels (c) and (d) can be compared with the equivalent results, obtained during model development, listed in Table C.1.




Figure C.5: Response of the Point Reyes forecast model to synthetic scenario NTSZ 30-39 (α = 25). Maximum sea surface elevation for (a) A grid, (b) B grid, (c) C grid. Sea surface elevation time series at the C grid warning point (d). Panels (c) and (d) can be compared with the equivalent results, obtained during model development, displayed in Figure 13.




Figure C.6: Response of the Point Reyes forecast model to the March 11 Tohoku tsunami. (a, b, and c) Maximum sea surface elevation for A, B and C grids. (d) Sea surface elevation time series at the C grid warning point. Panels (c) and (d) can be compared with the equivalent results, obtained during model development, displayed in Figure 17 and Figure 16a.

Scenario/Event	Source	Tsunami	ಶ	Maxima (cm)	ıa (cm)	Minima (cm)	a (cm)
Name	Zone	Source	(m)	SIFT	Development	SIFT	Development
			Mega-tsunami Scenarios	ni Scenarios			
KISZ 01-10	Kamchatka- Yap-Mariana- Izu-Bonin	A01-A10, B01-B10	25	354.127	354	-175.953	-177
KISZ 22-31	Kamchatka- Yap-Mariana- Izu-Bonin	A01-A10, B01-B10	25	250.248	251	-176.243	-176
ACSZ 56-65	Aleutian- Alaska- Cascadia	A01-A10, B01-B10	25	157.536	159	-158.122	-156
86-68 ZSSO	Central & South America	A01-A10, B01-B10	25	119.796	120	-141.905	-142
NTSZ 30-39	NewZealand- Kermadec- Tonga	A01-A10, B01-B10	25	401.252	402	-177.000	-177
			Historical Events	l Events			
Tohoku-2011	Kamchatka- Yap-Mariana-	4.66 b24 + 26.31 a26	4.66 b24 + 12.23 b25 + 26.31 a26 +21.27 b26 +	178.215	178	-148.389	-148
	Izu-Bonin	22.75 a27	22.75 a 27 + 4.98 b 27				

Table C.1: Table of maximum and minimum amplitudes (cm) at the Point Reyes, California warning point for synthetic and historical events tested using SIFT 3.2 and those obtained during development.

Glossary

- **Arrival time** The time when the first tsunami wave is observed at a particular location, typically given in local and/or universal time, but also commonly noted in minutes or hours relative to the time of the earthquake.
- **Bathymetry** The measurement of water depth of an undisturbed body of water.
- **Cascadia Subduction Zone** Fault that extends from Cape Mendocino in Northern California northward to mid-Vancouver Island Canada. The fault marks the convergence boundary where the Juan de Fuca tectonic plate is being subducted under the margin of the North America plate.
- **Current speed** The scalar rate of water motion measured as distance/time.
- **Current velocity** Movement of water expressed as a vector quantity. Velocity is the distance of movement per time coupled with direction of motion.
- **Digital Elevation Model (DEM)** A digital representation of bathymetry or topography based on regional survey data or satellite imagery. Data are arrays of regularly spaced elevations referenced to a map projection of the geographic coordinate system.
- **Epicenter** The point on the surface of the earth that is directly above the focus of an earthquake.
- **Focus** The point beneath the surface of the earth where a rupture or energy release occurs due to a buildup of stress or the movement of earth's tectonic plates relative to one another.
- **Inundation** The horizontal inland extent of land that a tsunami penetrates, generally measured perpendicularly to a shoreline.
- **Marigram** Tide gauge recording of wave level as a function of time at a particular location. The instrument used for recording is termed a marigraph.
- **Moment Magnitude** (M_W) The magnitude of an earthquake on a logarithmic scale in terms of the energy released. Moment magnitude is based on the size and characteristics of a fault rupture as determined from long-period seismic waves.
- **Method of Splitting Tsunamis (MOST)** A suite of numerical simulation codes used to provide estimates of the three processes of tsunami evolution: tsunami generation, propagation, and inundation.

- **Near-field** A particular location at which the earth's deformation due to energy release affects the modeling solution.
- **Propagation database** A basin-wide database of pre-computed water elevations and flow velocities at uniformly spaced grid points throughout the world oceans. Values are computed from tsunamis generated by earthquakes with a fault rupture at any one of discrete 100×50 km unit sources along worldwide subduction zones.
- **Runup** Vertical difference between the elevation of tsunami inundation and the sea level at the time of a tsunami. Runup is the elevation of the highest point of land inundated by a tsunami as measured relative to a stated datum, such as mean sea level.
- **Short-term Inundation Forecasting for Tsunamis (SIFT)** A tsunami forecast system that integrates tsunami observations in the deep-ocean with numerical models to provide an estimate of tsunami wave arrival and amplitude at specific coastal locations while a tsunami propagates across an ocean basin.
- **Subduction zone** A submarine region of the earth's crust at which two or more tectonic plates converge to cause one plate to sink under another, overriding plate. Subduction zones are regions of high seismic activity.
- **Synthetic event** Hypothetical events based on computer simulations or theory of possible or even likely future scenarios.
- **Tidal wave** Term frequently used incorrectly as a synonym for tsunami. A tsunami is unrelated to the predictable periodic rise and fall of sea level due to the gravitational attractions of the moon and sun: the tide.
- **Tide** The predictable rise and fall of a body of water (ocean, sea, bay, etc.) due to the gravitational attractions of the moon and sun.
- **Tide gauge** An instrument for measuring the rise and fall of a column of water over time at a particular location.
- **Tele–tsunami or distant tsunami or far–field tsunami** Most commonly, a tsunami originating from a source greater than 1000 km away from a particular location. In some contexts, a tele-tsunami is one that propagates through deep-ocean before reaching a particular location without regard to distance separation.
- **Travel time** The time it takes for a tsunami to travel from the generating source to a particular location.
- **tsunami** A Japanese term that literally translates to "harbor wave." Tsunamis are a series of long–period shallow water waves that are generated by the sudden displacement of water due to subsea disturbances such as earthquakes, submarine landslides, or volcanic eruptions. Less commonly, meteoric impact to the ocean or meteorological forcing can generate a tsunami.
- **Tsunami Hazard Assessment** A systematic investigation of seismically active regions of the world oceans to determine their potential tsunami impact at a particular location.

Numerical models are typically used to characterize tsunami generation, propagation, and inundation, and to quantify the risk posed to a particular community from tsunamis generated in each source region investigated.

- **Tsunami Propagation** The directional movement of a tsunami wave outward from the source of generation. The speed at which a tsunami propagates depends on the depth of the water column in which the wave is traveling. Tsunamis travel at a speed of 700 km/hr (450 mi/hr) over the average depth of 4000 m in the open deep Pacific Ocean.
- **Tsunami source** Location of tsunami origin, most typically an underwater earthquake epicenter. Tsunamis are also generated by submarine landslides, underwater volcanic eruptions, or, less commonly, by meteoric impact of the ocean.
- **Wave amplitude** The maximum vertical rise or drop of a column of water as measured from wave crest (peak) or trough to a defined mean water level state.
- **Wave crest or peak** The highest part of a wave or maximum rise above a defined mean water level state, such as mean lower low water.
- **Wave height** The vertical difference between the highest part of a specific wave (crest) and it's corresponding lowest point (trough).
- **Wavelength** The horizontal distance between two successive wave crests or troughs.
- **Wave period** The length of time between the passage of two successive wave crests or troughs as measured at a fixed location.
- **Wave trough** The lowest part of a wave or the maximum drop below a defined mean water level state, such as mean lower low water.